arXiv:1203.3236v2 [math.AG] 5 Apr 2012

On the birational section conjecture with local conditions

JAKOB STIX

Abstract — We prove that birational Galois sections for smooth curves over $\mathbb{Q}$ all have image in a decomposition group of a $\mathbb{Q}$-rational point if and only if certain associated 2-dimensional Galois representations are either reducible or form a compatible ℓ-adic system. This provides a birational variant endowed with an additional group theoretic condition of Grothendieck's section conjecture in anabelian geometry. We find here the first GL_2-type argument in anabelian geometry which usually only depends on prosolvable quotients.

As an aside we also obtain a strong approximation result for rational points on hyperbolic curves over $\mathbb{Q}$ or imaginary quadratic fields.

CONTENTS

1. Introduction 1
2. Galois sections with local conditions 3
3. Examples: semiabelian varieties 6
4. Finite support 9
5. Density of non-integral places 11
6. Cuspidal sections 18
7. On the birational section conjecture over $\mathbb{Q}$ with conditions 21
References 24

1. INTRODUCTION

1.1. **Motivation.** The section conjecture of anabelian geometry so far successfully refuses all attempts towards its proof: the conjecture in its original form remains open and is still a very hard problem despite some progress made over the last decade.

The section conjecture [Gr83] predicts that for a smooth hyperbolic curve X/k over a number field k every Galois section of the natural projection $\pi_1(X) \to \pi_1(\mathrm{Spec}(k))$ arises by functoriality from a rational point (possibly of the smooth completion), see [Sx11] for background and results.

When the emphasis is on diophantine applications, it clearly would be fascinating already to provide a description of the set of rational points in terms of sections with additional (group theoretic) conditions. Of course, we mean conditions different from the tautological condition of coming from a rational point. In this work, we impose three kinds of extra conditions:

(1) birational lifting: assume the existence of lifts to a section of $\pi_1(\mathrm{Spec}(K)) \to \pi_1(\mathrm{Spec}(k))$ where K is the function field of X, see Section 2.1,
(2) adelic: assume that the section locally belongs to an adelic point of X, see Section 2.2,
(3) ℓ-adic representations: if $k = \mathbb{Q}$, we assume that certain 2-dimensional Galois representations of Gal_k arising from families of elliptic curves over X are unramified outside a finite set of places and the place ℓ, see Section 5.3 and Theorem 31.

Both conditions (1) and (2) imply by Koenigsmann [Ko05] that the sections are Selmer, i.e., they come locally from points or are cuspidal. While (1) allows us to work birationally on the curve and is entirely in terms of groups and so is very much in the spirit of the original

---

Date: February 27, 2013.

conjecture and anabelian geometry, the second condition is rather strong as (2) imposes some kind of discreteness that we would rather deduce than invest in the theory. Nevertheless, the use of (2) is justified by the applications to strong approximation, see Theorem D below.

1.2. **Results.** Based on (1) and (2) we obtain a definite result for curves over totally real number fields or imaginary quadratic fields in Corollary 14.

Theorem A. *Let k be a totally real number field or an imaginary quadratic number field. Let X/k be a hyperbolic curve. Then the set of $\pi_1(X_{\bar{k}})$-conjugacy classes of birationally adelic sections of $\pi_1(X) \to \pi_1(\mathrm{Spec}(k))$ that are not cuspidal is in natural bijection with the set of k-rational points of X.*

We would like to emphasize that Theorem A does not rely on Faltings's Theorem, formerly known as the Mordell conjecture, which is important in light of initial hopes that the section conjecture would lead a new path towards that conjecture.

A weaker form of the second condition is the condition that sections are Selmer which in fact is implied by (1). At least up to a set S of places of density 0 we can then show that birationally liftable sections are either cuspidal or locally a rational point of X for $v \notin S$, see Corollary 23 and Theorem 28. It is remarkable that this part of the argument works over any number field k.

Theorem B. *Let $s : \mathrm{Gal}_k \to \pi_1(X)$ be a birationally liftable section of a hyperbolic curve X/k with smooth completion $\overline{X}$ over a number field k. Then the associated adele $\underline{a}(s) \in \overline{X}(\mathbb{A}_k)_\bullet$ has*

(1) either $a_v \in X(\mathfrak{o}_v)$ is integral for a set of places v of Dirichlet density 1,
(2) or the section s is cuspidal.

The proof of Theorem B uses in a crucial way the geometric monodromy of the Legendre family of elliptic curves. This leads to a new GL_2-type description of cuspidal sections in contrast to the characterization via weights of abelian nature due to Nakamura [Na90]. The use of the Legendre family has the flavour of the known reduction of the section conjecture for birationally liftable sections to the special case of $X = \mathbb{P}^1 - \{0, 1, \infty\}$ as observed for example in [EsHa08] Proposition 7.9. But in fact, we have to exploit many rational maps $X \dashrightarrow \mathbb{P}^1 - \{0, 1, \infty\}$ and so the line of thought is different.

In the case of hyperbolic curves over $\mathbb{Q}$, we offer the following version of a birational section conjecture with conditions, see Theorem 31. The limitation to $k = \mathbb{Q}$ comes from the use of Serre's modularity conjecture, now a Theorem by Khare and Wintenberger.

Theorem C. *Let $X/\mathbb{Q}$ be a hyperbolic curve. A birationally liftable section $s : \mathrm{Gal}_{\mathbb{Q}} \to \pi_1(X)$ comes from a rational point or is cuspidal, if and only if for every open $U \subseteq X$ with geometric point $\bar{u} \in U$, a lift $\tilde{s} : \mathrm{Gal}_{\mathbb{Q}} \to \pi_1(U, \bar{u})$, and every family E/U of elliptic curves with geometric fibre $E_{\bar{u}}$ over $\bar{u}$ the associated family of 2-dimensional representations*

$$\rho_{\tilde{s},E/U,\ell} = \rho_{E/U,\ell} \circ \tilde{s} \ : \ \mathrm{Gal}_{\mathbb{Q}} \to \mathrm{GL}(\mathrm{T}_\ell\, E_{\bar{u}})$$

has one of the following properties:

(i) There exists a finite set of places S independent of ℓ such that $\rho_{\tilde{s},E/U,\ell}$ is unramified outside ℓ and the places in S.
(ii) There is a character $\delta : \mathrm{Gal}_{\mathbb{Q}} \to \{\pm 1\}$ such that for all ℓ we have an exact sequence

$$0 \to \delta\varepsilon \to \rho_{\tilde{s},E/U,\ell} \to \delta \to 0$$

where ε is the ℓ-adic cyclotomic character.

1.3. **Outline.** We sketch the contents of the paper. Section 2 contains various notions of sections with (local) conditions. In Sections 3 and 4 we describe these notions for semi-abelian varieties and make use of Stoll's finite support result in descent theory. Here, as an aside, we obtain the following interesting result of strong approximation, see Corollary 13.

Theorem D. *Let X/k be a hyperbolic curve over either $k = \mathbb{Q}$ or k an imaginary quadratic number field such that $\mathcal{O}^*(X) \neq k^*$. Then the natural map*

$$X(k) \xrightarrow{\sim} X(\mathbb{A}_k)_\bullet^{\mathrm{f-desc}}$$

is a bijection from the set of rational points to the set of adelic points that survive any finite descent obstruction.

In Section 5 we start to draw conclusions for Selmer sections from the presence of non-constant families of elliptic curves. We prove a density theorem, Theorem 18, which relies on asymptotic of group theory in $\mathrm{GL}_2(\mathbb{F}_\ell)$ for $\ell \to \infty$. The theorem roughly says that the adele associated to a section behaves either, up to a set of density 0, like a rational point, or like a cuspidal section. This works over any number field. The precise statement concerning cuspidal sections is obtained in Section 6 by means of the geometric monodromy of the Legendre family.

In the final Section 7 we prove our characterization of birationally liftable sections that come from rational points in terms of 2-dimensional Galois representations.

Acknowledgements.

I would like to express my gratitude towards Olivier Wittenberg and Hélène Esnault for listening to the argument at an early stage when the state of affaires was still in total disorder. It is also my pleasure to thank Olivier Wittenberg for his comments on an earlier version of this manuscript. I thank David Harari for emphasizing the importance of strong approximation for semi-abelian varieties, and I am grateful to Leila Schneps for providing the explicit form of monodromy of for the Legendre family (actually in 2005 — a long time ago).

Notation and terminology.

A **hyperbolic curve** is a smooth relative curve $X \to S$ endowed with a smooth projective relative compactification $\overline{X} \to S$ with connected geometric fibres of constant genus such that $Y = \overline{X} \setminus X \to S$ is a finite étale relative divisor, and the fibrewise ℓ-adic Euler characteristic as a function $\bar{s} \mapsto \chi(X_{\bar{s}})$ on geometric points of S (where ℓ is invertible) is constant and negative.

For a number field k we denote its ring of integers by $\mathfrak{o}_k$, the completion at a place v of k is k_v with ring of integers $\mathfrak{o}_v$ if $v \nmid \infty$. The adele ring of k is denoted by $\mathbb{A}_k$. For a not necessarily finite set of places S of k we denote by $\mathfrak{o}_{k,S}$ the ring of S-integers (which are integral outside S).

2. Galois sections with local conditions

2.1. **Sections.** Let X/k be a geometrically connected irreducible and reduced variety with function field K. Let $\overline{K}$ be a fixed algebraic closure of K and set $\mathrm{Gal}_K = \mathrm{Gal}(\overline{K}/K)$. The algebraic closure of k contained in $\overline{K}$ will be denoted by $\bar{k}$ and $\mathrm{Gal}_k = \mathrm{Gal}(\bar{k}/k)$. The fundamental group of X/k forms an extension $\pi_1(X/k)$

$$1 \to \pi_1(X_{\bar{k}}) \to \pi_1(X) \to \mathrm{Gal}_k \to 1,$$

where the geometric generic point $\mathrm{Spec}(\overline{K}) \to X_{\bar{k}} \to X$ is the implicit base point. The space of sections of $\pi_1(X/k)$ up to conjugation by elements from $\pi_1(X_{\bar{k}})$ will be denoted by

$$\mathscr{S}_{\pi_1(X/k)}$$

and its birational analogue, the set of $\mathrm{Gal}_{K\bar{k}}$-conjugacy classes of sections of $\mathrm{Gal}_K \to \mathrm{Gal}_k$, will be denoted by

$$\mathscr{S}_{\pi_1(K/k)}.$$

By functoriality, to the point $a \in X(k)$ we associate a class of sections $s_a : \mathrm{Gal}_k \to \pi_1(X)$. This gives rise to the non-abelian profinite Kummer map $a \mapsto \kappa(a) = s_a$

$$\kappa : X(k) \to \mathscr{S}_{\pi_1(X/k)}.$$

2.1.1. *Birationally liftable sections.* Let $j : \operatorname{Spec} K \to X$ be the inclusion of the generic point which on π_1 induces the map

$$j_* : \mathrm{Gal}_K \to \pi_1(X),$$

a surjection if X is normal, and furthermore a map

$$j_* : \mathscr{S}_{\pi_1(K/k)} \to \mathscr{S}_{\pi_1(X/k)},$$

the image of which by definition is the set of **birationally liftable** sections

$$\mathscr{S}^{\mathrm{bir}}_{\pi_1(X/k)}.$$

We remark that the property *birationally liftable* used here is a priori stronger than the notion with the same name defined in [Sx11] §18.5.

2.1.2. *Neighbourhoods.* We recall the notion of a neighbourhood of a section $s : \mathrm{Gal}_k \to \pi_1(X)$. This is a finite étale cover $X' \to X$ together with a lift s' of the section, i.e., an open subgroup $H \subseteq \pi_1(X)$ containing the image of the section $s = s'$. The limit over all neighbourhoods yields a pro-étale cover

$$X_s = \varprojlim X' \to X$$

corresponding to $\pi_1(X_s) = s(\mathrm{Gal}_k) \subseteq \pi_1(X)$. It follows that a section s of $\pi_1(X/k)$ comes from a k-rational point of X if and only if $X_s(k)$ is nonempty. More precisely, we have $s = s_a$ if and only if

$$a \in \operatorname{im}\big(X_s(k) \to X(k)\big).$$

2.1.3. *Cuspidal section.* Let $X \subseteq \overline{X}$ be an open immersion with $\overline{X}$ smooth. Let $a \in \overline{X}(k)$ be a rational point not contained in X, and let w be a k-valuation of K with center in a on $\overline{X}$. Such valuations always exist due to the smoothness assumption, see [Sx12] Proposition 2. By general valuation theory, there is a split short exact sequence

$$1 \to I_w \to D_w \to \mathrm{Gal}_k \to 1$$

with I_w the inertia group and D_w the decomposition group of (a prolongation $\bar{w}$ to $\overline{K}$ of) the valuation w. Composing with the natural map

$$D_w \subseteq \mathrm{Gal}_K \to \pi_1(X)$$

leads to sections of $\pi_1(X/k)$ caused by the k-valuation w with center in $a \in (\overline{X} \setminus X)(k)$. These are by definition the **cuspidal sections** of $\pi_1(X/k)$, and these form naturally a subset

$$\mathscr{S}^{\mathrm{cusp}}_{\pi_1(X/k)} \subseteq \mathscr{S}^{\mathrm{bir}}_{\pi_1(X/k)}$$

of the birationally liftable sections. If a cuspidal section comes from a section of $D_w \to \mathrm{Gal}_k$, then we say that s belongs to the packet of cuspidal sections of the valuation w and that the cuspidal section is centered in a, the center of w on $\overline{X}$. Note that in this generality the packet in which a cuspidal section belongs is not necessarily unique.

Let now X be a smooth curve. For a section $s : \mathrm{Gal}_k \to \pi_1(X)$ we consider the pro-(finite branched) cover

$$\overline{X}_s = \varprojlim_{X'} \overline{X'}$$

where $X' \to X$ ranges through neighbourhoods of s and $\overline{X'}$ is the normalization of $\overline{X}$ in $X' \to X$. Then s is cuspidal and centered in a if and only if

$$a \in \operatorname{im}\big(\overline{X}_s(k) \to \overline{X}(k)\big) \setminus X(k).$$

2.2. **Local conditions.** Form now on, let k be a number field and let $k \hookrightarrow k_v$ denote the completion at a place v of k. In analogy to Selmer groups in the Galois cohomology of abelian varieties which classify torsors that become locally trivial because they possess local points everywhere, we would like to consider Selmer sections as sections that at least locally belong to a rational point.

2.2.1. *Base change of sections.* Since we are now in characteristic 0 we have a base change map for sections, see [Sx11] §3.2, with respect to an extension of fields k'/k:

$$\mathscr{S}_{\pi_1(X/k)} \to \mathscr{S}_{\pi_1(X\times_k k'/k')} =: \mathscr{S}_{\pi_1(X/k)}(k')$$
$$s \mapsto s \otimes k'$$

Note that in general base change does not exist for birational sections, only if k'/k is algebraic:

$$\mathscr{S}_{\pi_1(K/k)} \to \mathscr{S}_{\pi_1(Kk'/k')},$$

because, if k'/k is transcendental, then $\mathrm{Gal}_{Kk'} \to \mathrm{Gal}_K \times_{\mathrm{Gal}_k} \mathrm{Gal}_{k'}$ not an isomorphism.

2.2.2. *Selmer sections.* In case of a number field k and v a place and $k' = k_v$ we write $s\otimes k_v = s_v$. Then, we furthermore say that a section $s : \mathrm{Gal}_k \to \pi_1(X)$ is a **Selmer section** if for every place v of k the localisation s_v is cuspidal or belongs to a rational point in $X(k_v)$. The subset of all Selmer sections we denote by

$$\mathscr{S}^{\mathrm{Sel}}_{\pi_1(X/k)} \subseteq \mathscr{S}_{\pi_1(X/k)}.$$

For a hyperbolic curve X/k with smooth projective completion $\overline{X}$, the adelic point of $\overline{X}$ associated to a Selmer section is unique and leads to a well defined map

$$\underline{a} \,:\, \mathscr{S}^{\mathrm{Sel}}_{\pi_1(X/k)} \to \overline{X}(\mathbb{A}_k)_{\bullet}$$
$$s \mapsto \underline{a}(s) = (a_v)_v$$

with defining property $s_v = s_{a_v}$. Here $\overline{X}(\mathbb{A}_k)_{\bullet}$ denotes the set of modified adelic points of X where the component at an infinite place v is given by $\pi_0(X(k_v))$.

It follows form [HSx10] Theorem 11 that the map $\underline{a}$ takes values in the set $\overline{X}(\mathbb{A}_k)_{\bullet}^{\mathrm{f-desc}}$ of adelic points that survive all finite descent obstructions, so that we obtain a map

$$\underline{a} \,:\, \mathscr{S}^{\mathrm{Sel}}_{\pi_1(X/k)} \to \overline{X}(\mathbb{A}_k)_{\bullet}^{\mathrm{f-desc}}. \tag{2.1}$$

Moreover, if $X = \overline{X}$ is a smooth projective curve of genus at least 1, then by [HSx10] Theorem 11 the map (2.1) is even surjective.

2.2.3. *Birationally liftable sections are Selmer.* We recall Koenigsmann's Lemma [Ko05] Proposition 2.4 (a) and its consequence [Ko05] Corollary 2.6.

Proposition 1. *Let X/k be a hyperbolic curve over a number field k with smooth projective completion $X \subseteq \overline{X}$ and function field K/k.*

(1) If $\pi_1(X/k)$ admits a birationally liftable section, then $\overline{X}(k_v) \neq \emptyset$ for every completion k_v.
(2) Any birationally liftable section of $\pi_1(X/k)$ is a Selmer section:

$$\mathscr{S}^{\mathrm{bir}}_{\pi_1(X/k)} \subseteq \mathscr{S}^{\mathrm{Sel}}_{\pi_1(X/k)}.$$

Proof: (1) If $\pi_1(X/k)$ admits a birationally liftable section, then $\mathrm{Gal}_K \to \mathrm{Gal}_k$ splits and (1) is [Ko05] Corollary 2.6.

(2) Now we apply (1) to all neighbourhoods of s. Since $\overline{X}_{s_v} = \overline{X}_s \times_k k_v$ we find

$$\overline{X}_s(k_v) = \varprojlim_{X'} \overline{X'}(k_v) \neq \emptyset$$

as the X' range over neighbourhoods of s, because this is a projective limit of nonempty compact sets. Consequently, with

$$a_v \in \mathrm{im}(\overline{X}_s(k_v) \to \overline{X}(k_v))$$

we have $s_v = s_{a_v}$, either as the section associated to $a_v \in X(k_v)$, or as a cuspidal section associated to a choice of splitting of

$$D_{a_v} \to \mathrm{Gal}_{k_v}$$

where D_{a_v} is the decomposition subgroup of $a_v \in \overline{X} \setminus X$. In any case, the section s is Selmer. $\square$

2.2.4. *Adelic sections.* Let X/k be a hyperbolic curve. An **adelic section** is a Selmer section $s : \mathrm{Gal}_k \to \pi_1(X)$ such that the associated adele of the smooth completion $\underline{a}(s) \in \overline{X}(\mathbb{A}_k)_\bullet$ lies in the subset of adelic points $X(\mathbb{A}_k)_\bullet$ of X. The set of all adelic sections will be denoted by

$$\mathscr{S}^{\mathrm{adelic}}_{\pi_1(X/k)},$$

in particular, we obtain a map

$$\underline{a} \, : \, \mathscr{S}^{\mathrm{adelic}}_{\pi_1(X/k)} \twoheadrightarrow X(\mathbb{A}_k)_\bullet^{\mathrm{f-desc}}$$

that is surjective due to [HSx10] Theorem 11, see also [Sx11] Theorem 144.

2.2.5. *Birationally adelic sections.* Let X/k be a hyperbolic curve. A **birationally adelic section** is a section $s : \mathrm{Gal}_k \to \pi_1(X)$ that is birationally lifting to a section $\mathrm{Gal}_k \to \mathrm{Gal}_K$ such that for every open $U \subseteq X$ the induced section of $\pi_1(U/k)$ is either adelic for U/k or cuspidal. The set of all birationally adelic sections is denoted by

$$\mathscr{S}^{\mathrm{ba}}_{\pi_1(X/k)}.$$

To summarize, we have defined the following hierarchy of sections with (local) conditions:

$$\begin{tikzcd}
\mathscr{S}^{\mathrm{cusp}}_{\pi_1(X/k)} \amalg X(k) \arrow[r, phantom, "\subseteq"] \arrow[dd, two heads] & \mathscr{S}^{\mathrm{ba}}_{\pi_1(X/k)} \arrow[r, phantom, "\subseteq"] \arrow[d, phantom, sloped, "\subseteq"] & \mathscr{S}^{\mathrm{bir}}_{\pi_1(X/k)} \arrow[d, phantom, sloped, "\subseteq"] & \\
 & \mathscr{S}^{\mathrm{adelic}}_{\pi_1(X/k)} \amalg \mathscr{S}^{\mathrm{cusp}}_{\pi_1(X/k)} \arrow[r, phantom, "\subseteq"] \arrow[d, two heads, "\underline{a}"] & \mathscr{S}^{\mathrm{Selmer}}_{\pi_1(X/k)} \arrow[r, phantom, "\subseteq"] \arrow[d, two heads, "\underline{a}"] & \mathscr{S}_{\pi_1(X/k)} \\
\overline{X}(k) \arrow[r, hook] & X(\mathbb{A}_k)_\bullet^{\mathrm{f-desc}} \amalg (\overline{X} \setminus X)(k) \arrow[r, phantom, "\subseteq"] & \overline{X}(\mathbb{A}_k)_\bullet^{\mathrm{f-desc}}. &
\end{tikzcd}$$

Question 2. It is natural to ask whether birationally liftable sections are in fact either adelic or cuspidal, whereby, if the answer is always yes, the hierarchy simplifies to

$$\mathscr{S}^{\mathrm{cusp}}_{\pi_1(X/k)} \amalg X(k) \subseteq \mathscr{S}^{\mathrm{ba}}_{\pi_1(X/k)} = \mathscr{S}^{\mathrm{bir}}_{\pi_1(X/k)} \subseteq \mathscr{S}^{\mathrm{adelic}}_{\pi_1(X/k)} \amalg \mathscr{S}^{\mathrm{cusp}}_{\pi_1(X/k)} \subseteq \mathscr{S}^{\mathrm{Selmer}}_{\pi_1(X/k)} \subseteq \mathscr{S}_{\pi_1(X/k)}.$$

3. EXAMPLES: SEMIABELIAN VARIETIES

Let B/k be a semi-abelian variety over a number field k and let

$$0 \to T \to B \to A \to 0$$

be the associated extension of an abelian variety A/k by a torus T/k. By continuous Kummer theory we obtain an exact sequence

$$0 \to \widehat{B(k)} \to \mathscr{S}_{\pi_1(B/k)} \to \mathrm{Hom}\left(\mathbb{Q}/\mathbb{Z}, \mathrm{Div}(\mathrm{H}^1(k,B))\right) \to 0$$

where $\widehat{M} := \varprojlim_n M/nM$ and $\mathrm{Div}(M)$ denotes the maximal divisible subgroup of M. By Hilbert's Theorem 90, the group $\mathrm{H}^1(k,T)$ is annihilated by the degree of an extension that splits T. Hence the projection $\mathrm{pr} : B \to A$ induces an injective map

$$\mathrm{pr}_* \, : \, \mathrm{Div}(\mathrm{H}^1(k,B)) \hookrightarrow \mathrm{Div}(\mathrm{H}^1(k,A)).$$

A section $s : \mathrm{Gal}_k \to \pi_1(B)$ is Selmer if and only if $\mathrm{pr}_*(s)_v$ comes from a point in $A(k_v) = \widehat{A(k_v)}$ for every place v of k, so that the following is exact:

$$0 \to \widehat{B(k)} \to \mathscr{S}^{\mathrm{Selmer}}_{\pi_1(B/k)} \to \mathrm{Hom}\big(\mathbb{Q}/\mathbb{Z}, \mathrm{Div}(Ш(A/k))\big) \to 0. \tag{3.1}$$

It follows that the group of Selmer sections is nothing but the continuous Selmer group

$$\mathscr{S}^{\mathrm{Selmer}}_{\pi_1(B/k)} = \mathrm{H}^1_{\mathrm{Sel}}\big(k, \prod_\ell \mathrm{T}_\ell(B)\big)$$

with coefficients in the Tate module $\prod_\ell \mathrm{T}_\ell(B)$ of B.

Proposition 3. *Let B/k be a semi-abelian variety such that $Ш(A/k)$ is finite for the maximal abelian variety quotient $B \to A$. Then the natural map*

$$B(k) \to \mathscr{S}^{\mathrm{adelic}}_{\pi_1(B/k)}$$

is bijective if and only if $B(k) \subset B(\mathbb{A}_k)_\bullet$ is a discrete subset.

Proof: The finiteness of the Tate–Shafarevich group $Ш(A/k)$ implies by the above that

$$\widehat{B(k)} = \mathscr{S}^{\mathrm{Selmer}}_{\pi_1(B/k)}.$$

Since the natural map

$$\widehat{B(k)} \to \prod_{v\nmid\infty} \widehat{B(k_v)} \times \prod_{v|\infty} \pi_0(B(k_v))$$

is injective, we can view adelic sections as the intersection

$$\widehat{B(k)} \cap B(\mathbb{A}_k)_\bullet = \mathscr{S}^{\mathrm{adelic}}_{\pi_1(B/k)}$$

inside that product space. Intersecting with $\widehat{B(k)}$ imposes on an adelic point in $B(\mathbb{A}_k)_\bullet$ that it has to survive all finite descent obstructions. Since for a semi-abelian variety the fundamental group is abelian and torsion free, the finite descent obstruction to adelic points agrees with the Brauer–Manin obstruction. We deduce by strong approximation [Ha08] Theorem 4 and our assumption of $Ш(A/k)$ being finite that

$$\overline{B(k)} = \widehat{B(k)} \cap B(\mathbb{A}_k)_\bullet^{\mathrm{f-desc}} = \mathscr{S}^{\mathrm{adelic}}_{\pi_1(B/k)}$$

where the topological closure of the rational points is taken in $B(\mathbb{A}_k)_\bullet$ with respect to the natural topology. □

Here are three special cases for Proposition 3 that we will exploit in the sequel. These examples admit more elementary proofs which are given for the convenience of the reader. The first example treats $\mathbb{G}_m$.

Proposition 4. *Let k be $\mathbb{Q}$ or a quadratic imaginary number field. Then $\mathbb{G}_m$ over $\mathrm{Spec}(k)$ has*

$$k^* = \mathbb{G}_m(k) = \mathscr{S}^{\mathrm{adelic}}_{\pi_1(\mathbb{G}_m/k)}.$$

Proof: The diagonal map $\widehat{k^*} \hookrightarrow \prod_v \widehat{k_v^*}$ is injective, see [NSW08] Theorem 9.1.11(2). The intersection

$$\widehat{k^*} \cap \mathbb{G}_m(\mathbb{A}_k)_\bullet$$

inside the product contains k^*. By finiteness of the class number we can fix a finite set of places S of k containing all infinite places S_∞ with $\mathrm{Pic}(\mathfrak{o}_{k,S}) = 1$. Then we can use elements from k^* to move the support of the divisor of any $(a_v) \in \mathbb{G}_m(\mathbb{A}_k)_\bullet$ into S and thus

$$\widehat{k^*} \cap \mathbb{G}_m(\mathbb{A}_k)_\bullet = k^* \cdot \left(\widehat{k^*} \cap \prod_{v\notin S} \mathfrak{o}_v^* \times \prod_{v\in S\setminus S_\infty} k_v^* \times \prod_{v\in S_\infty} \pi_0(k_v^*) \right)$$

$$= k^* \cdot \left(\widehat{\mathfrak{o}^*_{k,S}} \cap \prod_{v \notin S} \mathfrak{o}_v^* \times \prod_{v \in S \setminus S_\infty} k_v^* \times \prod_{v \in S_\infty} \pi_0(k_v^*) \right)$$

$$= k^* \cdot \ker(v \otimes \hat{\mathbb{Z}} : \mathfrak{o}_{k,S} \otimes \hat{\mathbb{Z}} \to \bigoplus_{v \in S \setminus S_\infty} \hat{\mathbb{Z}}/\mathbb{Z}) = k^* \cdot \widehat{\mathfrak{o}_k^*}.$$

The number fields under consideration here are exactly those with $\mathfrak{o}_k^*$ finite. Hence we conclude that

$$k^* = k^* \cdot \widehat{\mathfrak{o}_k^*} = \mathscr{S}^{\text{adelic}}_{\pi_1(\mathbb{G}_m/k)}$$

and the proof is complete. □

The second example is concerned with a nontrivial torus of rank 1.

Proposition 5. *Let F be a totally real number field and let E/F be a quadratic extension that is totally imaginary. Then, for the norm 1-torus*

$$T = T_{E/F} = \ker(N : R_{E|F}\mathbb{G}_m \to \mathbb{G}_m)$$

over F we have

$$T(F) = \ker(N : E^* \to F^*) = \mathscr{S}^{\text{adelic}}_{\pi_1(T/F)}.$$

Proof: We need to compute $\widehat{T(F)} \cap T(\mathbb{A}_F)_\bullet$ which certainly injects by restriction into

$$\ker\left(N : \widehat{E^*} \cap \mathbb{G}_m(\mathbb{A}_E)_\bullet \to \widehat{F^*} \cap \mathbb{G}_m(\mathbb{A}_F)_\bullet\right) = \ker\left(N : E^* \cdot \widehat{\mathfrak{o}_E^*} \to F^* \cdot \widehat{\mathfrak{o}_F^*}\right),$$

where we have used the general computation of Proposition 4. By Dirichtlet's Unit Theorem the map $N : \mathfrak{o}_E^* \to \mathfrak{o}_F^*$ is an isomorphism up to torsion. This is preserved under profinite completion, and furthermore, the map

$$N : \widehat{\mathfrak{o}_E^*}/\mathfrak{o}_E^* \to \widehat{\mathfrak{o}_F^*}/\mathfrak{o}_F^*$$

is an isomorphism (of uniquely divisible groups isomorphic to $(\hat{\mathbb{Z}}/\mathbb{Z})^{d-1}$ where $d = [F : \mathbb{Q}]$). It follows from the snake lemma applied to

$$\begin{tikzcd}
0 \arrow[r] & E^* \arrow[r] \arrow[d, "N"] & E^* \cdot \widehat{\mathfrak{o}_E^*} \arrow[r] \arrow[d, "N"] & \widehat{\mathfrak{o}_E^*}/\mathfrak{o}_E^* \arrow[r] \arrow[d, "N", "\cong"'] & 0 \\
0 \arrow[r] & F^* \arrow[r] & F^* \cdot \widehat{\mathfrak{o}_F^*} \arrow[r] & \widehat{\mathfrak{o}_F^*}/\mathfrak{o}_F^* \arrow[r] & 0
\end{tikzcd}$$

that the natural map

$$T(F) = \ker(N : E^* \to F^*) \xrightarrow{\sim} \ker\left(N : E^* \cdot \widehat{\mathfrak{o}_E^*} \to F^* \cdot \widehat{\mathfrak{o}_F^*}\right)$$

is an isomorphism. This completes the proof. □

The third example treats abelian varieties and was observed by Stoll.

Proposition 6. *Let A/k be an abelian variety such that $A(k)$ and $Ш(A/k)$ are finite. Then we have naturally*

$$A(k) = \mathscr{S}^{\text{adelic}}_{\pi_1(A/k)} = \mathscr{S}^{\text{Selmer}}_{\pi_1(A/k)}.$$

Proof: Selmer and adelic sections clearly agree for all proper varieties. The proof now follows directly from (3.1). □

Example 7. Examples for abelian varieties over $\mathbb{Q}$ that satisfy the rather restrictive assumptions of Proposition 6 are classically found as follows. A modular abelian variety is an abelian variety $A/\mathbb{Q}$ which occurs as a quotient $J_0(N) \to A$ of a modular Jacobian for suitable level N. The Birch and Swinnerton-Dyer conjecture predicts that $A(\mathbb{Q})$ and $Ш(A/\mathbb{Q})$ are finite as soon as $A/\mathbb{Q}$ has trivial analytic rank, i.e., its critical L-value $L(A/\mathbb{Q}, 1) \neq 0$ does not vanish. By Mazur's work on the Eisenstein ideal we know that indeed for every prime $N = 11$ or $N \geq 17$ there are modular abelian varieties $J_0(N) \to A$ of trivial analytic rank and Kolyvagin and Logachëv show in [KL89] [KL92] that indeed in these cases $A(\mathbb{Q})$ and $Ш(A/\mathbb{Q})$ are finite.

4. FINITE SUPPORT

4.1. **The support of a Selmer section.** Let X/k be a hyperbolic curve over the number field k with smooth projective completion $\overline{X}$. The **support** of a Selmer section $s : \mathrm{Gal}_k \to \pi_1(X)$ is defined as the Zariski-closed subscheme of $\overline{X}$

$$Z(s) = \overline{\bigcup_v \mathrm{im}(a_v : \mathrm{Spec}(k_v) \to \overline{X})}$$

where $\underline{a}(s) = (a_v) \in \overline{X}$ is the adelic point associated to the Selmer section. We say that a Selmer section s has **finite support** if $Z(s)$ is finite over k.

4.2. **Finite support and sections.** On adelic points with finite support we have the following important result due to Stoll.

Theorem 8 (Stoll [St07] Theorem 8.2)**.** *Let Z be a proper closed subscheme of a smooth projective curve $\overline{X}$ of genus at least 1 over a number field k. Then the diagonal map*

$$Z(k) \xrightarrow{\sim} \Big\{(a_v) \in \overline{X}(\mathbb{A}_k)_\bullet^{\mathrm{f-desc}} \ ; \ a_v \in Z(k_v) \textit{ for a set of places } v \textit{ of density } 1\Big\}$$

is a bijection.

Stoll's result is even stronger in that the adelic point only needs to survive finite descent obstructions with respect to abelian groups. For our applicatioins to sections the difference does not matter.

Corollary 9. *Let X/k be a hyperbolic curve over a number field. Then the image of the map*

$$X(k) \to \mathscr{S}^{\mathrm{adelic}}_{\pi_1(X/k)} \subseteq \mathscr{S}^{\mathrm{Selmer}}_{\pi_1(X/k)} \setminus \mathscr{S}^{\mathrm{cusp}}_{\pi_1(X/k)}$$

consists precisely of the non-cuspidal Selmer sections with finite support.

Proof: For $a \in X(k)$ the support of s_a is the subscheme $\{a\} \hookrightarrow \overline{X}$. It remains to conclude the converse: if the support $Z(s)$ of the Selmer section s is finite and s is not cuspidal, then s belongs to some rational point of X. If we pass to a neighbourhood $h : X' \to X$ of the section s with lift s', then

$$Z(s') \subseteq h^{-1}(Z(s)),$$

so that the property of having finite support is preserved. We may therefore without loss of generality assume that the smooth completion $\overline{X}$ of X has genus ≥ 1. Then by Theorem 8

$$\underline{a}(s) \in Z(s)(\mathbb{A}_k)_\bullet \cap \overline{X}(\mathbb{A}_k)_\bullet^{\mathrm{f-desc}} = Z(k)$$

and so in the limit over all neighbourhoods of s

$$X_s(k) \supseteq \varprojlim_{(X',s')} Z(s')(k) \neq \emptyset.$$

The limit argument recalled in Section 2.1 implies that $s = s_a$ for any $a \in \mathrm{im}\big(\overline{X}_s(k) \to \overline{X}(k)\big)$. Since we assumed s not to be cuspidal, we may even deduce that $a \in X(k)$. ☐

Remark 10. It is extremely remarkable that the version of the limit argument performed in the course of the proof of Corollary 9 does not rely on Faltings' Theorem, formerly known as the Mordell conjecture. Usually, the passage from existence of a rational point on a curve over a number field to knowing that a given section arises from a rational point requires the limit $\varprojlim_{X'} X'(k)$ over all neighbourhoods of s to be a limit of compact sets. This compactness is usually guaranteed by Faltings' Theorem dashing the original hope that a proof of the section conjecture provides an alternative approach to this basic diophantine finiteness. Here, our a priori knowledge that the rational point will lie in Z allows to deal with a subset of this limit which is trivially compact.

Corollary 11. *Let X/k be a hyperbolic curve that admits a nonconstant map $f : X \to B$ to a semi-abelian variety B/k such that*

(1) for the abelian variety quotient $B \to A$ we have $Ш(A/k)$ is finite, and
(2) $B(k) \subset B(\mathbb{A}_k)_\bullet$ is a discrete subset.

Then all adelic sections of $\pi_1(X/k)$ have finite support and in particular the natural maps

$$X(k) \xrightarrow{\sim} \mathscr{S}^{\text{adelic}}_{\pi_1(X/k)} \xrightarrow{\sim} X(\mathbb{A}_k)^{\text{f-desc}}_\bullet$$

given by κ and $\underline{a}$ are bijective.

Proof: We first treat κ. By Corollary 9 it suffices to show that adelic sections $s : \mathrm{Gal}_k \to \pi_1(X)$ have finite support. Clearly

$$Z(s) \subseteq f^{-1}(Z(f_*s))$$

and so it suffices to consider the adelic section $f_*s : \mathrm{Gal}_k \to \pi_1(B)$. By Proposition 3 and assumptions (i) and (ii) we know that $f_*s = s_b$ for some $b \in B(k)$. Hence $Z(s)$ is contained in the finite subscheme $f^{-1}(b) \subset X$ which concludes the proof that $X(k) = \mathscr{S}^{\text{adelic}}_{\pi_1(X/k)}$.

The map $X(k) \hookrightarrow X(\mathbb{A}_k)^{\text{f-desc}}_\bullet$ is injective and $\underline{a} : \mathscr{S}^{\text{adelic}}_{\pi_1(X/k)} \twoheadrightarrow X(\mathbb{A}_k)^{\text{f-desc}}_\bullet$ is surjective by [HSx10] Theorem 11. In view of $X(k) = \mathscr{S}^{\text{adelic}}_{\pi_1(X/k)}$ we coclude that also $\underline{a}$ must be bijective which finishes the proof. ☐

Remark 12. The argument of Corollary 11 was given by Stoll in the case of $B = A$ being an abelian variety, see [St07] Theorem 8.6.

Corollary 13. *Let k be $\mathbb{Q}$ or an imaginary quadratic number field. Let X/k be a hyperbolic curve with a global nonconstant unit $f \in \mathcal{O}_X^* \setminus k^*$. Then the natural maps*

$$X(k) \xrightarrow{\sim} \mathscr{S}^{\text{adelic}}_{\pi_1(X/k)} \xrightarrow{\sim} X(\mathbb{A}_k)^{\text{f-desc}}_\bullet$$

given by κ and $\underline{a}$ are bijective.

Proof: The nonconstant unit defines a nonconstant map $f : X \to \mathbb{G}_m$. Corollary 11 together with Proposition 4 conclude the proof. ☐

Corollary 14. *Let k be a totally real number field or an imaginary quadratic number field. Let X/k be a hyperbolic curve. Then the natural map*

$$\mathscr{S}^{\text{cusp}}_{\pi_1(X/k)} \amalg X(k) \xrightarrow{\sim} \mathscr{S}^{\text{ba}}_{\pi_1(X/k)}$$

is a bijection.

Proof: We have to show that birationally adelic sections $s : \mathrm{Gal}_k \to \pi_1(X)$ have finite support in order to conclude with Corollary 9. For this we may replace X by a nonempty open $U \subset X$. If $k/\mathbb{Q}$ is imaginary quadratic we choose U such that we have a nonconstant global unit on U and conclude with Corollary 13.

It remains to treat the case of a totally real number field k. Let $a \in k^*$ be a totally negative element and set $k' = k(\sqrt{a})$ with $\mathrm{Gal}(k'/k)$ generated by σ. We denote the norm-1 torus of the totally imaginary quadratic extension k'/k by

$$T = \ker(N : R_{k'|k}\mathbb{G}_m \to \mathbb{G}_m).$$

We denote the function field of X by $k(X)$. A rational map $f : X \dashrightarrow T$ corresponds to an element $f \in k(X) \otimes_k k'$ of norm

$$N(f) = \sigma(f)f = \mathbf{1}.$$

We take $\alpha \in k' \setminus k$ and a nonconstant element $g \in k(X)^* \setminus k^*$ and set

$$f = \sigma(g + \alpha)/(g + \alpha) = (g + \sigma(\alpha))/(g + \alpha).$$

Then clearly $N(f) = \mathbf{1}$ and f is nonconstant. We pick $U \subset X$ small enough such that $f : U \to T$ is an algebraic map. Now Corollary 11 together with Proposition 5 applied to adelic sections of $\pi_1(U/k)$ and the map $f : U \to T$ conclude the proof. $\square$

A result in the same spirit but using abelian varieties instead of tori was proven by Stoll [St07] Theorem 8.6 and Remark 8.9.

Corollary 15 (Stoll [St07])**.** *Let X/k be a hyperbolic curve over a number field k with a nontrivial map $f : X \to A$ for some abelian variety A/k with finite $A(k)$ and finite $Ш(A/k)$. Then every Selmer section $s : \mathrm{Gal}_k \to \pi_1(X)$ is cuspidal or $s = s_a$ for some $a \in X(k)$, so the natural maps provide identifications*

$$\mathscr{S}^{\mathrm{cusp}}_{\pi_1(X/k)} \amalg X(k) \xrightarrow{\sim} \mathscr{S}^{\mathrm{ba}}_{\pi_1(X/k)} = \mathscr{S}^{\mathrm{bir}}_{\pi_1(X/k)} = \mathscr{S}^{\mathrm{adelic}}_{\pi_1(X/k)} \amalg \mathscr{S}^{\mathrm{cusp}}_{\pi_1(X/k)} = \mathscr{S}^{\mathrm{Selmer}}_{\pi_1(X/k)}$$

and

$$\overline{X}(\mathbb{A}_k)_\bullet^{\mathrm{f-desc}} = \overline{X}(k).$$

Proof: Since we work with a map $X \to A$ to an abelian variety, the argument of Corollary 11 applies to all Selmer sections. The proof of the corollary thus follows from Proposition 6. $\square$

5. Density of non-integral places

We now discuss to what extent a Selmer section misses to be an adelic or cuspidal section.

5.1. **Types of adelic points.** Let X/k be a smooth, geometrically connected curve over a number field k and let $\overline{X}$ be its smooth projective completion. Let $U \subseteq \mathrm{Spec}(\mathfrak{o}_k)$ be a dense open such that $X \subseteq \overline{X}$ has good reduction $\mathscr{X} \subseteq \overline{\mathscr{X}}$ over U in the sense of open curves, i.e., $\mathscr{X}$ is an open in the smooth, projective $\overline{\mathscr{X}} \to U$ and the boundary divisor $\overline{\mathscr{X}} \setminus \mathscr{X}$ is relatively étale over U. For an adelic point

$$\underline{a} = (a_v) \in \overline{X}(\mathbb{A}_k)_\bullet$$

we define a partition of all places of k with respect to U and the models $\mathscr{X} \subseteq \overline{\mathscr{X}}$ as follows.

integral : An integral place for $\underline{a}$ is a place $v \in U$ such that the closure of $\{a_v\}$ in $\overline{\mathscr{X}} \times_U \mathfrak{o}_v$ is contained in $\mathscr{X} \times_U \mathfrak{o}_v$.

degenerate : A degenerate place for $\underline{a}$ is a place $v \in U$ such that $a_v \in X(k_v)$ and the closure of $\{a_v\}$ in $\overline{\mathscr{X}} \times_U \mathfrak{o}_v$ meets the boundary $(\overline{\mathscr{X}} \setminus \mathscr{X}) \times_U \mathfrak{o}_v$.

cuspidal : A cuspidal place for $\underline{a}$ is a place $v \in U$ such that $a_v \in (\overline{X} \setminus X)(k_v)$.

bad : A bad place for $\underline{a}$ is a place $v \notin U$. In particular, all infinite places are bad by definition.

Definition 16. With the notation as above, we say that $(a_v) \in \overline{X}(\mathbb{A}_k)_\bullet$ is **asymptotically integral** if the intersection number with the boundary divisor $\overline{\mathscr{X}} \setminus \mathscr{X}$ of the closure of $\{a_v\}$ for degenerate places v tends to 0 in $\hat{\mathbb{Z}}$, i.e., for every $n \geq 1$ there are only finitely many degenerate places v where n does not divide this intersection number.

Remark 17. (1) Since we will be interested in assertions on the finiteness or the Dirichlet density of the partition sets, the choice of U and the models $\mathscr{X} \subseteq \overline{\mathscr{X}}$ are irrelevant.

(2) The subset $X(\mathbb{A}_k)_\bullet \subseteq \overline{X}(\mathbb{A}_k)_\bullet$ contains precisely those adelic points for which all but finitely many places are integral, no place is cuspidal and the components at $v \notin U$ lie in $X(k_v)$.

5.2. **Families of elliptic curves and ℓ-adic representations.** Let X/k be a geometrically connected variety with geometric point $\bar{x} \in X$. We assume that there is a family of elliptic curves

$$E \to X$$

and consider, for every $\ell \neq \mathrm{char}(k)$ the corresponding ℓ-adic 2-dimensional representation

$$\rho_{E/X,\ell} : \pi_1(X, \bar{x}) \to \mathrm{GL}(\mathrm{T}_\ell(E_{\bar{x}}))$$

where $E_{\bar{x}}$ is the geometric fibre of E/X in $\bar{x}$. The Weil-pairing induces a canonical isomorphism

$$\det(\rho_{E/X,\ell}) = \varepsilon \circ \mathrm{pr}_*$$

where ε is the corresponding ℓ-adic cyclotomic charater and $\mathrm{pr}_* : \pi_1(X,\bar{x}) \to \mathrm{Gal}_k$ is induced by the projection map.

To any section $s : \mathrm{Gal}_k \to \pi_1(X,\bar{x})$ we can thus associate a family of ℓ-adic representations

$$\rho_{s,E/X,\ell} = \rho_{E/X,\ell} \circ s \ : \ \mathrm{Gal}_k \to \mathrm{GL}(\mathrm{T}_\ell(E_{\bar{x}}))$$

with cyclotomic determinant

$$\det(\rho_{s,E/X,\ell}) = \varepsilon.$$

By naturality of the construction, if $s = s_a$ for $a \in X(k)$, then $\rho_{s,E/X,\ell}$ is nothing but the Galois representation $\rho_{E_a/k,\ell}$ on $\mathrm{T}_\ell(E_a)$ for the fibre E_a of E/X in a. If the family is constant $E = X \times E_0$, then clearly $\rho_{s,E/X,\ell}$ is independent of s and agrees with $\rho_{E_0/k,\ell}$.

Recall that k is a number field. Strictly speaking, the family $\{\rho_{s,E/X,\ell}\}_\ell$ does not deserve to be called an ℓ-adic representation, because

(i) we lack a conductor N such that $\rho_{s,E/X,\ell}$ is unramified for all places $v \nmid \ell \cdot N$,
(ii) we lack integrality of the characteristic polynomials of Frobenius,
(iii) and we lack independence of ℓ for the characteristic polynomials of Frobenius.

This changes for Selmer sections, well almost.

5.3. **Almost an ℓ-adic representation.** Let now X/k be a smooth, geometrically connected curve, and assume that the family E/X has bad semistable reduction along every point of $\overline{X} \setminus X$. Let $s : \mathrm{Gal}_k \to \pi_1(X)$ be a Selmer section with associated adelic point $\underline{a}(s) = (a_v)$ of the smooth projective completion $\overline{X}$. Let $D_v = \mathrm{Gal}_{k_v} \subset \mathrm{Gal}_k$ be the decomposition subgroup of the place v, and let $I_v \subset D_v$ be the inertia subgroup. We discuss the local behaviour

$$\rho_{s,E/X,\ell}|_{D_v} : \mathrm{Gal}_{k_v} \to \mathrm{GL}(\mathrm{T}_\ell(E_{\bar{x}}))$$

in terms of the type of v with respect to $\underline{a}$. As $\mathscr{X} \to U$ with $U \subseteq \mathrm{Spec}(\mathfrak{o}_k)$ we take a model such that the family E/X has good reduction over $\mathscr{X}$ and bad semistable reduction along $\overline{\mathscr{X}} \setminus \mathscr{X}$. Note that $\overline{X}$ is a surface and that semistable reduction ceases to make sense only in a set of codimension 2, hence a finite set that we may assume to be empty by shrinking U.

integral: For an integral v the local representation belongs to the elliptic curve E_v/k_v which is the fibre of E/X in $a_v \in X(k_v)$ and has good reduction over $\mathrm{Spec}(\mathfrak{o}_v)$. It follows that

(i) the representation $\rho_{s,E/X,\ell}|_{D_v}$ is unramified for $\ell \neq \mathrm{char}(\mathbb{F}_v)$ with $\mathbb{F}_v$ being the residue field of v,
(ii) the characteristic polynomial of Frobenius is integral

$$\det(\mathbf{1} - \mathrm{Frob}_v T | \rho_{s,E/X,\ell}) = \mathbf{1} - a_v T + N(v) T^2 \in \mathbb{Z}[T]$$

(here $N(v) = \#\mathbb{F}_v$ is the norm of v),
(iii) and the trace of Frobenius $a_v \in \mathbb{Z}$ is independent of ℓ with $|a_v| \leq 2\sqrt{N(v)}$.

degenerate: For a degenerate v the local representation belongs to the elliptic curve E_v/k_v which is the fibre of E/X in $a_v \in X(k_v)$ and has bad semistable reduction over $\mathrm{Spec}(\mathfrak{o}_v)$. It follows that

(i) there is a quadratic unramified character $\delta_v : \mathrm{Gal}_{k_v} \to \{\pm 1\}$ such that for all $\ell \neq \mathrm{char}(\mathbb{F}_v)$ in a suitable basis

$$\rho_{s,E/X,\ell}|_{D_v} \sim \begin{pmatrix} \delta\varepsilon & \psi \\ & \delta \end{pmatrix} \tag{5.1}$$

where $\psi|_{I_v} = m_v \cdot t_\ell$ is a multiple of the tame ℓ-adic character with $m_v > 0$ integral and independent of ℓ,

(ii) the characteristic polynomial of Frobenius still makes sense (computed on the unramified semisimplification) and is integral

$$\begin{aligned}\det(\mathbf{1} - \mathrm{Frob}_v T|\rho_{s,E/X,\ell}) &= \mathbf{1} - a_v T + N(v)T^2 \\ &= (\mathbf{1} - \delta_v(\mathrm{Frob}_v)T)(\mathbf{1} - \delta_v(\mathrm{Frob}_v)N(v)T) \in \mathbb{Z}[T],\end{aligned}$$

(iii) and the trace of Frobenius $a_v \in \mathbb{Z}$ is independent of ℓ with

$$a_v = \delta_v(\mathrm{Frob}_v)(N(v) + 1).$$

The character δ describes the 1-dimensional torus in the special fibre of the Néron model of E_v over $\mathrm{Spec}(\mathfrak{o}_v)$. Concerning the claim on ψ we may pass to an unramified extension k'_v/k_v such that E_v attains split multiplicative reduction and therefore admits a Tate uniformization $E_v \times_{k_v} k'_v = \mathbb{G}_m/q_v^{\mathbb{Z}}$, with $q_v \in k'_v$. The cocycle $\psi|_{I_v}$ is the Kummer cocycle associated to q_v and thus agrees with the $m_v = v(q_v) > 0$ multiple of the tame ℓ-adic character.

cuspidal: For a cuspidal v the local section s_v is cuspidal at a_v and thus factors over the decomposition subgroup of a_v in $\pi_1(X \otimes k_v/k_v)$: the absolute Galois group of the fraction field of $\hat{\mathcal{O}}_{\overline{X}\otimes k_v, a_v}$ that is noncanonically isomorphic to $k_v((z))$. It follows that the image of the representation $\rho_{s,E/X,\ell}|_{D_v}$ is contained in the image of the representation of the corresponding Tate elliptic curve which is the fibre $E \times_X \mathrm{Spec}\, k((z))$. We thus can say the same thing about $\rho_{s,E/X,\ell}|_{D_v}$ for $v \nmid \ell$ as for degenerate places v except that we do know nothing on ψ:

(i) There is a quadratic unramified character $\delta_v : \mathrm{Gal}_{k_v} \to \{\pm 1\}$ such that for all $\ell \neq \mathrm{char}(\mathbb{F}_v)$ in a suitable basis

$$\rho_{s,E/X,\ell}|_{D_v} \sim \begin{pmatrix} \delta\varepsilon & * \\ & \delta \end{pmatrix},$$

(ii) the characteristic polynomial of Frobenius still makes sense and is integral

$$\begin{aligned}\det(\mathbf{1} - \mathrm{Frob}_v T|\rho_{s,E/X,\ell}) &= \mathbf{1} - a_v T + N(v)T^2 \\ &= (\mathbf{1} - \delta_v(\mathrm{Frob}_v)T)(\mathbf{1} - \delta_v(\mathrm{Frob}_v)N(v)T) \in \mathbb{Z}[T],\end{aligned}$$

(iii) and the trace of Frobenius $a_v \in \mathbb{Z}$ is independent of ℓ with

$$a_v = \delta_v(\mathrm{Frob}_v)(N(v) + 1).$$

bad: For the finitely many bad v we say nothing.

For a Selmer section s as above and $\rho = \{\rho_{s,E/X,\ell}\}$ we define the **trace of Frobenius**

$$a_v(\rho) = \mathrm{tr}\big(\rho_{s,E/X,\ell}(\mathrm{Frob}_v)|\,\mathrm{T}_\ell(E_{\bar{x}})\big) \in \mathbb{Z}$$

for any $\ell \neq \mathrm{char}(\mathbb{F}_v)$ and any v that is not bad. By the above discussion this number is indeed a well defined integer.

5.4. **A dichotomoy.** As a consequence of the Chebotarev Density Theorem we will now prove the following result.

Theorem 18. *Let X/k be a hyperbolic curve over a number field with smooth completion $\overline{X}$. Let E/X be a family of elliptic curves with bad and not even potentially good reduction over $\overline{X} \setminus X$. Let $s : \mathrm{Gal}_k \to \pi_1(X)$ be a Selmer section with adelic point $\underline{a}(s) = (a_v) \in \overline{X}(\mathbb{A}_k)_\bullet$. Then $\underline{a}(s)$ is asymptotically integral with respect to X, and exactly one of the following occurs.*

(1) Either the set

$$\{\textit{places } v \textit{ such that } a_v \textit{ is integral with respect to } X\}$$

has Dirichlet density 1,

(2) or the family of ℓ-adic representations $\rho_{s,E/X,\ell}$ factors in a suitable basis through

$$\begin{pmatrix} \delta\varepsilon & * \\ & \delta \end{pmatrix}$$

with a quadratic character $\delta : \mathrm{Gal}_k \to \{\pm 1\}$ that is independent of ℓ and ramified at most in the bad places. Moreover, all but finitely many places are cuspidal or degenerate. The remaining places are bad.

Proof: We may pass to a neighbourhood $h : X' \to X$ of the section s. For every $y' \in \overline{X'} \setminus X'$ above $y \in \overline{X} \setminus X$, the semistable reduction theorem implies that h^*E/X' has semistable reduction at y' if the ramification index $e_{y'/y}$ is divisible by an integer that only depends on the degeneration of E in y. Since X is hyperbolic, among the neighbourhoods of the section s we find universal ramification along $\overline{X} \setminus X$. Hence we may and will assume from the beginning that the family E/X has bad semistable reduction outside X.

Let us first address the claim on asymptotic integrality. Let ℓ be a prime number and let $r \geq 1$. We have to show divisibility by ℓ^r of the intersection number d_v of the closure of $\{a_v\}$ with the boundary for almost all degenerate $v \nmid \ell$.

Let $m_v = v(q_v) > 0$ be the valuation of the local Tate parameter q_v at a degenerate place v for the elliptic curve E_v as in Section 5.3 above. Being essentially a finite quotient of a global Galois group, the mod ℓ^r reduction of the representation $\rho_{s,E/X,\ell}$ is unramified for allmost all v. If $v \nmid \ell$ is degenerate and unramified in the mod ℓ^r representation, then $\ell^r \mid m_v$ by the description of ψ in (5.1).

Let now t be a local parameter on $\overline{\mathscr{X}} \times_U \mathfrak{o}_v$ for the boundary component of $(\overline{\mathscr{X}} \setminus \mathscr{X}) \times_U \mathfrak{o}_v$ that intersects with a_v. The j-function on $\mathscr{X}$ induced by the family E/X (more precisely its extension to $\mathscr{X}$) has a pole along $\{t = 0\}$ of some order e and thus $t^e \sim j^{-1}$ differ by a v-adic unit. Moreover, the local Tate parameter q_v has $v(q_v) = -v(j(a_v))$. This leads to

$$d_v = v(t(a_v)) = -\frac{1}{e} v(j(a_v)) = \frac{1}{e} v(q_v) = \frac{m_v}{e}.$$

Since only finitely many e can occur, we conclude asymptotic integrality for $\underline{a}(s)$.

We now address the claimed dichotomy. Let $G_\ell \subseteq \mathrm{GL}_2(\mathbb{F}_\ell)$ be the image of the mod ℓ reduction of $\rho_{s,E/X,\ell}$. And let $M_\ell \subseteq G_\ell$ be the subset of elements with split characteristic polynomial and at least one eigenvalue ± 1 modulo ℓ. The Frob_v for v cuspidal or degenerate are contained in M_ℓ so that their Dirichlet density is bounded above by

$$\frac{\# M_\ell}{\# G_\ell}.$$

We will show that this ratio can become arbitrarily small for ℓ ranging over all prime numbers or otherwise $\rho_{s,E/X,\ell}$ has the exceptional form (2).

Since the determinant of our representation is the cyclotomic character, we may assume by working with $\ell \gg 0$ that

$$\det \ : \ G_\ell \twoheadrightarrow \mathbb{F}_\ell^*$$

is surjective.

Let PG_ℓ be the image of G_ℓ in $\mathrm{PGL}_2(\mathbb{F}_\ell)$. Then the classification of subgroups $G \subseteq \mathrm{GL}_2(\mathbb{F}_\ell)$ with $\det(G) = \mathbb{F}_\ell^*$ (for $\ell \gg 0$) says that we can have either of the following cases:

(i) The mod ℓ representation is reducible, i.e.,

$$G_\ell \subseteq \begin{pmatrix} * & * \\ & * \end{pmatrix},$$

(ii) $G_\ell = \mathrm{GL}_2(\mathbb{F}_\ell)$,

(iii) G_ℓ is contained in the normalizer of a split torus but not in the torus and $\ell \nmid \# G_\ell$,

(iv) G_ℓ is contained in the normalizer of a non-split torus and $\ell \nmid \# G_\ell$,

(v) PG_ℓ is either A_4, S_4, or S_5 and $\ell \nmid \#G_\ell$.

By Lemmata 19–22 below we are done unless case (i) occurs for all but finitely many places. We therefore now assume that all mod ℓ representations are reducible for $\ell \gg 0$.

Let $\chi_i : G_\ell \to \mathbb{F}_\ell^*$ for $i = 1, 2$ be the projection onto the two diagonal entries. We denote by $H_i \subseteq \mathbb{F}_\ell^*$ the subgroup generated by $\chi_i(M_\ell)$. If the index

$$(\chi(G_\ell) : H_i)$$

is unbounded for ℓ ranging over all prime numbers, then $\#M_\ell/\#G_\ell \leq \#H_i/\#\chi_i(G_\ell)$ becomes arbitrarily small and we are done. We therefore assume that the two indices are bounded.

Let $\overline{G}_\ell$ denote the image of G_ℓ under

$$\mathrm{pr} = (\chi_1, \chi_2) : \begin{pmatrix} * & * \\ & * \end{pmatrix} \to \mathbb{F}_\ell^* \times \mathbb{F}_\ell^*$$

and set $\overline{M}_\ell = \mathrm{pr}(M_\ell)$. We have

$$(H_1)^2 \times (H_2)^2 \subseteq \overline{G}_\ell$$

and therefore the estimate

$$\frac{\#M_\ell}{\#G_\ell} \leq \frac{\#\overline{M}_\ell}{\#\overline{G}_\ell} \leq \frac{2\#H_1 + 2\#H_2}{1/4 \cdot \#H_1 \cdot \#H_2} = \frac{8}{\#H_1} + \frac{8}{\#H_2}$$

so that we can conclude the claim of the theorem if both $\#H_i$ are unbounded when ℓ ranges over all prime numbers.

It remains to discuss the case, where $\min_{i=1,2}\{\#\chi_i(G_\ell)\}$ remains bounded when ℓ tends to infinity. In this case there is an $m \in \mathbb{N}$ independent of ℓ and v such that there is a $\zeta \in \mu_m$ depending on ℓ, v such that

$$a_v(\rho) = \mathrm{tr}(\mathrm{Frob}_v \,|\rho_{s,E/X,\ell}) \equiv \zeta + \zeta^{-1}N(v) \mod \ell.$$

Now there are only finitely many $\zeta \in \mu_m$ and so for a each fixed v there must be one ζ that is good for infinitely many ℓ. With that ζ we find

$$a_v(\rho) = \zeta + \zeta^{-1}N(v)$$

so that ζ satisfies a nontrivial quadratic relation over $\mathbb{Z}$. If $[\mathbb{Q}(\zeta) : \mathbb{Q}] = 2$, then this must be irreducible, and $N(v) = N(\zeta) = 1$ which is absurd. Therefore $\zeta \in \mathbb{Q}$ and thus $\zeta = \pm 1$. We conclude that

$$a_v(\rho) = \pm(N(v) + 1)$$

for every finite place v. This contradicts the Hasse–Weil bound $|a_v(\rho)| \leq 2\sqrt{N(v)}$ in case v were integral. We deduce that all but finitely many places are cuspidal or degenerate and the remaining places are bad (the same argument shows that the two scenarios (1) and (2) of the theorem cannot hold simultaneously).

By approximating an arbitrary element $\sigma \in \mathrm{Gal}_k$ by Frobenius elements at places which are unramified in the mod ℓ^n reduction of $\rho_{s,E/X,\ell}$ we deduce that in $\mathbb{Z}_\ell$.

$$\mathrm{tr}(\rho_{s,E/X,\ell}(\sigma)) = \pm(\varepsilon(\sigma) + 1).$$

Here the sign is independent of ℓ since we can approximate the mod ℓ^n reduction for two different ℓ simultaneously by Frobenius elements and there the sign is independent of ℓ.

Let $\Gamma_\ell \subset \mathrm{GL}(\mathrm{T}_\ell(E_{\bar{x}}))$ be the image of the representation $\rho_{s,E/X,\ell}$. Then Γ_ℓ is a closed ℓ-adic analytic group and by [La65] contains an open normal subgroup

$$\Gamma_\ell^0 \lhd \Gamma_\ell$$

that consists of squares from Γ_ℓ. Let k_ℓ be the finite extension of k corresponding to the finite quotient $\mathrm{Gal}_k \twoheadrightarrow \Gamma_\ell/\Gamma_\ell^0$. Then for $\sigma \in \mathrm{Gal}_{k_\ell}$ we find $\gamma \in \Gamma_\ell$ with

$$\rho_{s,E/X,\ell}(\sigma) = \gamma^2.$$

Using the identity for $A \in GL_2$

$$\mathrm{tr}(A^2) = \mathrm{tr}(A)^2 - 2\det(A),$$

we compute

$$\begin{aligned} \mathrm{tr}(\rho_{s,E/X,\ell}(\sigma)) &= \mathrm{tr}(\gamma^2) = (\mathrm{tr}(\gamma))^2 - 2\det(\gamma) \\ &= (\pm(\varepsilon(\gamma)+1))^2 - 2\varepsilon(\gamma) = \varepsilon(\gamma^2) + 1 = \varepsilon(\sigma) + 1. \end{aligned}$$

It follows that the semisimplification of $\rho_{s,E/X,\ell}|\mathrm{Gal}_{k_\ell}$ agrees with $\varepsilon \oplus \mathbf{1}$. Since $\varepsilon \neq \mathbf{1}$ and the trivial representation is preserved by automorphisms we conclude that $\rho_{s,E/X,\ell}$ itself is also reducible. The semisimplification must be $\delta\varepsilon \oplus \delta^{-1}$ with a character δ of finite order. For the Frobenius elements, we find values of these characters

$$\{\pm 1, \pm N(v)\} = \{\delta(\mathrm{Frob}_v)N(v), \delta^{-1}(\mathrm{Frob}_v)\}.$$

As $\pm N(v)$ is never torsion in $\mathbb{Z}_\ell^*$ we must have $\delta(\mathrm{Frob}_v) = \pm 1$ and therefore δ is a quadratic character

$$\delta : \mathrm{Gal}_k \to \{\pm 1\} \subset \mathbb{Z}_\ell^*.$$

Moreover, δ is independent of ℓ, since it is determined by the signs in $a_v(\rho) = \pm(N(v)+1)$ which are independent of ℓ. Furthermore, by comparing with the local form at cuspidal or degenerate places, we see that δ can be ramified at most at the bad places.

It remains to exclude that $\rho_{s,E/X,\ell}$ has the form

$$\begin{pmatrix} \delta & * \\ & \delta\varepsilon \end{pmatrix}$$

in a suitable basis without being the direct sum $\delta \oplus \delta\varepsilon$. Assume that this happens. Then no place can be degenerate since inertia would then act nontrivially unipotently and thereby uniquely determine the fixed $\mathbb{Z}_\ell$-line. However, the description of the local representations then says that the character associated to this submodule must be $\delta\varepsilon$, a contradiction. If now all but finitely many places are cuspidal, we conclude that the adelic point $\underline{a}(s)$ has finite support in $\overline{X} \setminus X$. This allows to use Corollary 9 (or better its proof) to deduce that s is cuspidal. But then it follows from the known structure (as recalled above for cuspidal places) of the Galois representation associated to Tate elliptic curves that $\rho_{s,E/X,\ell}$ has the shape of (2). This finally finishes the proof of the theorem. □

5.5. **Asymptotics in subgroups of $\mathrm{GL}_2(\mathbb{F}_\ell)$.** We now provide the Lemmas needed in the proof of Theorem 18.

Lemma 19. *If $G_\ell = \mathrm{GL}_2(\mathbb{F}_\ell)$, then*

$$\#M_\ell / \#G_\ell \leq 2/(\ell - 1).$$

Proof: The possible Jordan normal forms for elements in M_ℓ are

$$\begin{pmatrix} 1 & \\ & 1 \end{pmatrix}, \begin{pmatrix} -1 & \\ & -1 \end{pmatrix}, \begin{pmatrix} a & \\ & 1 \end{pmatrix}, \begin{pmatrix} b & \\ & -1 \end{pmatrix}, \begin{pmatrix} 1 & 1 \\ & 1 \end{pmatrix}, \begin{pmatrix} -1 & 1 \\ & -1 \end{pmatrix}$$

with $a, b \in \mathbb{F}_\ell^*$ and, to make the list disjoint, the condition $a \neq 1$ and $b \neq \pm 1$. We now have to sum up the reciprocals of the size of the respective centralizer. This leads to

$$\begin{aligned} \frac{\#M_\ell}{\#G_\ell} &= \frac{2}{(\ell^2-1)(\ell^2-\ell)} + \frac{2\ell-5}{(\ell-1)^2} + \frac{2}{\ell(\ell-1)} \\ &= \frac{2\ell}{(\ell^2-1)(\ell-1)} + \frac{2\ell-5}{(\ell-1)^2} \leq \frac{2\ell-3}{(\ell-1)^2} \leq \frac{2}{\ell-1}. \end{aligned}$$

□

Lemma 20. *If $PG_\ell = A_4$, S_4, or A_5 and $\det(G_\ell) = \mathbb{F}_\ell^*$, then*

$$\#M_\ell / \#G_\ell \leq 60/(\ell - 1).$$

Proof: We consider an element $A \in M_\ell$ with eigenvalues $a, 1$ or $-a, -1$. The order of the image of A in PG_ℓ is in both cases the order of $a \in \mathbb{F}_\ell^*$. Since the order of an element in A_4, S_4, or A_5 divides 60 we conclude that a must lie in the 60-torsion of $\mathbb{F}_\ell^*$.

Since $\det(A) = a$ we conclude that $\det(M_\ell)$ is also contained in the 60-torsion of $\mathbb{F}_\ell^*$. The estimate

$$\frac{\#M_\ell}{\#G_\ell} \leq \frac{\#\det(M_\ell)}{\ell - 1} \leq \frac{60}{\ell - 1}$$

finishes the proof. □

Lemma 21. *If G_ℓ is contained in the normalizer of a non-split torus, $\ell \nmid \#G_\ell$ and $\det(G_\ell) = \mathbb{F}_\ell^*$, then*

$$\#M_\ell / \#G_\ell \leq 2/(\ell - 1).$$

Proof: The normalizer of a nonsplit torus has the form $\mathbb{F}_{\ell^2}^* \rtimes \mathrm{Gal}(\mathbb{F}_{\ell^2}/\mathbb{F}_\ell)$. We consider an element $A \in M_\ell$ with eigenvalues $a, 1$ or $-a, -1$. Then A^2 is contained in the non-split torus with eigenvalues $a^2, 1$. The eigenvalues of $\lambda \in \mathbb{F}_{\ell^2}$ are $\lambda, \bar{\lambda}$. We deduce that necessarily $a^2 = 1$, and the Jordan normal form of A is one of the following

$$\begin{pmatrix} 1 & \\ & 1 \end{pmatrix}, \begin{pmatrix} -1 & \\ & -1 \end{pmatrix}, \begin{pmatrix} 1 & \\ & -1 \end{pmatrix}.$$

Therefore $\det(M_\ell)$ is contained in the 2-torsion of $\mathbb{F}_\ell^2$ and the estimate

$$\frac{\#M_\ell}{\#G_\ell} \leq \frac{\#\det(M_\ell)}{\ell - 1} \leq \frac{2}{\ell - 1}$$

finishes the proof. □

Lemma 22. *If G_ℓ is contained in the normalizer of a split torus but not in a torus, $\ell \nmid \#G_\ell$ and $\det(G_\ell) = \mathbb{F}_\ell^*$, then*

$$\#M_\ell / \#G_\ell \leq 6/\sqrt{(\ell - 1)}.$$

Proof: The normalizer of the split torus is

$$(\mathbb{F}_\ell^* \times \mathbb{F}_\ell^*) \rtimes \langle \begin{pmatrix} & 1 \\ 1 & \end{pmatrix} \rangle = \left\{ \begin{pmatrix} a & \\ & b \end{pmatrix}, \begin{pmatrix} & b \\ a & \end{pmatrix} \ ; \ a, b \in \mathbb{F}_\ell^* \right\}.$$

For $A \in M_\ell$ not in the torus, we have $A = \begin{pmatrix} & b \\ a & \end{pmatrix}$ with characteristic polynomial $X^2 - ab$. As ±1 must be a root we see that $ab = 1$. Moreover, two such elements differ by an element of

$$D := G_\ell \cap \mathrm{SL}_2(\mathbb{F}_\ell),$$

and their eigenvalues are ±1.

Let $H \subseteq \mathbb{F}_\ell^*$ be the subgroup generated by all eigenvalues of elements from M_ℓ. Then for every $a \in H^2$ we have

$$\begin{pmatrix} a & \\ & 1 \end{pmatrix} \in G_\ell.$$

Therefore we have $H^2 \times H^2 \subseteq G_\ell$ and the product map $D \times (H^2 \times \{\mathbf{1}\}) \hookrightarrow G_\ell$ is injective. Since H is cyclic, we obtain the estimates

$$(\#H)^2 \leq 4\#G_\ell \qquad \text{and} \qquad \#H \cdot \#D \leq 2\#G_\ell.$$

Now we simply count the elements in M_ℓ by counting those of the form

$$\begin{pmatrix} a & \\ & \pm 1 \end{pmatrix}, \begin{pmatrix} \pm 1 & \\ & a \end{pmatrix}, \begin{pmatrix} & a^{-1} \\ a & \end{pmatrix}$$

so that

$$\#M_\ell \leq 4 \cdot \#H + \#D.$$

As $\det(M_\ell) \subseteq \pm H$ has size $\#\det(M_\ell) \leq 2\#H$, the estimate

$$\left(\frac{\#M_\ell}{\#G_\ell}\right)^2 \leq \frac{4\cdot\#H + \#D}{\#G_\ell} \cdot \frac{\#\det(M_\ell)}{\ell - 1}$$
$$\leq \frac{8(\#H)^2 + 2\#D\cdot\#H}{\#G_\ell(\ell-1)} \leq \frac{36}{\ell-1}$$

finishes the proof. □

Corollary 23. *Let $s : \mathrm{Gal}_k \to \pi_1(X)$ be a birationally liftable section of a hyperbolic curve with smooth completion $\overline{X}$ over a number field k. Then the associated adele $\underline{a}(s) \in \overline{X}(\mathbb{A}_k)_\bullet$ has*

(1) either $a_v \in X(\mathfrak{o}_v)$ is integral for a set of places v of Dirichlet density 1,

(2) or all but finitely many places v are cuspidal or degenerate with respect to $X \subset \overline{X}$.

Proof: The open subsets $U = \beta^{-1}(\mathbb{P}^1 - \{0, 1, \infty\}) \subseteq X$ for finite maps $\beta : \overline{X} \to \mathbb{P}^1$ which map $\overline{X} \setminus X$ to $\{0, 1, \infty\}$ form a basis of the topology of X (even with β étale over $\mathbb{P}^1 - \{0, 1, \infty\}$ due to an improved version of Belyi's Theorem by Mochizuki [Mz04] Corollary 1.1). The Legendre family of elliptic curves

$$E_\lambda = \{Y^2 = X(X-1)(X-\lambda)\} \;\to\; \mathbb{P}^1 - \{0, 1, \infty\}$$

has bad reduction exactly in 0, 1, and ∞. Thus we can apply Theorem 18 to a lift $\mathrm{Gal}_k \to \pi_1(U)$ of s and the pullback family $\beta^* E_\lambda \to U$. It follows that either $a_v \in U(\mathfrak{o}_v)$ is integral for a set of places of density 1, in which case we are done, or secondly that all but finitely many places are cuspidal or degenerate with respect to $U \subseteq \overline{X}$.

We may therefore assume that we are in the second case for all U as above. Let X be covered by $U_1, \ldots, U_n$ for open subsets U_i as above (in fact two such sets suffice). Let $\mathscr{Y}_i$ be the Zariski closure of $X \setminus U_i$ in a suitable common model. The intersection $\bigcap_i \mathscr{Y}_i$ is finite. Thus a_v is actually cuspidal or degenerate also for allmost all places v with respect to $X \subset \overline{X}$. □

6. Cuspidal sections

6.1. **The geometric monodromy of the Legendre family.** For the finer analysis in Theorem 28 below we have to understand the geometric monodromy representation of the Legendre family

$$\rho_{\mathrm{Leg}} = \rho_{E_\lambda/\mathbb{P}^1-\{0,1,\infty\}} \;:\; \pi_1(\mathbb{P}^1_{\overline{\mathbb{Q}}} - \{0,1,\infty\}, \overrightarrow{01}) \to \mathrm{GL}_2(\mathbb{Z}_\ell).$$

In fact the 2-adic representaion turns out to be crucial. Since we are in characteristic 0, this is nothing but the profinite/pro-2 completion of the topological monodromy on the period lattice of the family. The topological fundamental group

$$\pi_1^{\mathrm{top}}(\mathbb{P}^1(\mathbb{C}) - \{0,1,\infty\}, \overrightarrow{01}) = \langle x, y, z | xyz = 1\rangle$$

is freely generated by an infinitesimal counterclockwise loop x around 0 starting and ending at the tangential base point $\overrightarrow{01}$, and by the path y which is the image of x under $\lambda \mapsto 1 - \lambda$ conjugated by the path from $\overrightarrow{01}$ to $\overrightarrow{10}$ along the real interval $[0,1]$. The path $z = (xy)^{-1}$ turns out to be a an infinitesimal loop around ∞ conjugated by a path form $\overrightarrow{01}$ to a tangential base point at ∞. In particular, this topological presentation reveals representatives for the inertia groups at the cusps 0, 1, and ∞, namely $I_0 = \langle x\rangle$, $I_1 = \langle y\rangle$, and $I_\infty = \langle z\rangle$.

Using this description of $\pi_1^{\mathrm{top}}(\mathbb{P}^1(\mathbb{C}) - \{0,1,\infty\}, \overrightarrow{01})$ the following is classical:

Lemma 24. *For a suitable basis, the topological monodromy representation*

$$\rho_{\mathrm{Leg}}^{\mathrm{top}} \;:\; \pi_1^{\mathrm{top}}(\mathbb{P}^1(\mathbb{C}) - \{0,1,\infty\}, \overrightarrow{01}) \to \mathrm{GL}_2(\mathbb{Z})$$

for the Legendre family maps the generators as follows:

$$x \mapsto \begin{pmatrix} 1 & 2 \\ & 1 \end{pmatrix}, \qquad y \mapsto \begin{pmatrix} 1 & \\ -2 & 1 \end{pmatrix}, \qquad z \mapsto \begin{pmatrix} 1 & -2 \\ 2 & -3 \end{pmatrix}.$$

Proof: Well known computation of the monodromy of the Picard-Fuchs equation associated to the family of elliptic curves. ☐

Remark 25. Note that the explicit formulae of Lemma 24 allow to conclude that inertia at 0 and 1 acts unipotently, while inertia at ∞ acts quasi-unipotently with Jordan normal form

$$\begin{pmatrix} -1 & 1 \\ & -1 \end{pmatrix}$$

which corresponds to the fact that the Legendre family is semistable at $0, 1$ and additive at ∞.

6.2. **Unipotent subgroups up to conjugation.** Let $U \subseteq \mathrm{GL}_2(\mathbb{Z}_\ell)$ be a nontrivial unipotent subgroup. Then

$$L_U = \ker(\mathbf{1} - U) \subset \mathbb{Z}_\ell \times \mathbb{Z}_\ell$$

is a free and cotorsion free submodule of rank 1 and as such defines an element $L_U \in \mathbb{P}^1(\mathbb{Z}_\ell)$. Conversely, to such a line $L \in \mathbb{P}^1(\mathbb{Z}_\ell)$ we can define the unipotent subgroup

$$U(L) = \{A \in \mathrm{GL}_2(\mathbb{Z}_\ell) \ ; \ A|_L = \mathrm{id}_L \ \text{ and } \ \det(A) = 1\}.$$

Clearly, $U \subseteq U(L_U)$ and unipotent subgroups of the form U_L are maximal among unipotent subgroups with respect to inclusion.

Lemma 26. *Every nontrivial unipotent subgroup U of $\mathrm{GL}_2(\mathbb{Z}_\ell)$ is contained in a unique maximal unipotent subgroup, namely $U(L_U)$. The map $U \mapsto L_U$ defines a bijection*

$$\{\textit{maximal unipotent subgroups of } \mathrm{GL}_2(\mathbb{Z}_\ell)\} \longleftrightarrow \mathbb{P}^1(\mathbb{Z}_\ell).$$

Proof: The map $L \mapsto U(L)$ is the inverse map. ☐

6.3. **Recognizing cusps via unipotent subgroups.** It follows from Lemma 26 that $\mathrm{GL}_2(\mathbb{Z}_\ell)$ acts transitively by conjugation on the set of its maximal unipotent subgroups. However, this changes if we consider only the conjugation action by a suitable subgroup.

Lemma 27. *The maximal unipotent subgroups U_0, U_1, and U_∞ of $\mathrm{GL}_2(\mathbb{Z}_2)$ containing the images of (the square of) inertia in the 2-adic geometric monodromy representation of the Legendre family*

$$\rho_{\mathrm{Leg}}(I_0), \rho_{\mathrm{Leg}}(I_1), \ \textit{and respectively } \rho_{\mathrm{Leg}}((I_\infty)^2)$$

are mutually not conjugate under the image $\rho_{\mathrm{Leg}}\big(\pi_1(\mathbb{P}^1_{\bar{\mathbb{Q}}} - \{0,1,\infty\}, \overrightarrow{01})\big)$.

Proof: The Legendre family has trivial 2-torsion as is reflected by $\rho_{\mathrm{Leg}}^{\mathrm{top}}$ mapping to the subgroup $\Gamma(2) \subset \mathrm{SL}_2(\mathbb{Z})$ of elements $\equiv \mathbf{1} \bmod 2$. This is inherited by the profinite completion ρ_{Leg}. We deduce that conjugation by elements from $\mathrm{im}(\rho_{\mathrm{Leg}})$ only moves maximal unipotent subgroups within the fibre of the mod 2 reduction

$$\mathbb{P}^1(\mathbb{Z}_2) \twoheadrightarrow \mathbb{P}^1(\mathbb{F}_2) = \{0, 1, \infty\}.$$

Now x fixes $\binom{1}{0}$, and y fixes $\binom{0}{1}$ and z^2 fixes $\binom{1}{1}$, so that U_0, U_1, and U_∞ map to three different elements in $\mathbb{P}^1(\mathbb{F}_2)$. ☐

6.4. **Enough families of elliptic curves.** We are now in a position to treat Selmer sections that behave like cuspidal sections with respect to the dichotomy of Theorem 18.

Theorem 28. *Let $s : \mathrm{Gal}_k \to \pi_1(X)$ be a birational lifting section of a hyperbolic curve X over a number field k with smooth completion $\overline{X}$. If the associated adele $\underline{a}(s) = (a_v)$ is cuspidal or degenerate with respect to $X \subseteq \overline{X}$ at all but finitely many places of k, then s is a cuspidal section of $\pi_1(X/k)$.*

Proof: Let $Y = \overline{X} \setminus X$ be the complement. Since the assumptions are inherited by neighbourhoods, it suffices to show that $Y(k)$ is nonempty and then apply the limit argument as recalled in Section 2.1.

We consider the following map defined on almost all places of k:

$$v \mapsto y_v \in Y$$

where we assign to a cuspidal or degenerate place v of k the closed point y_v of Y such that the closure of a_v and y_v in a model over $\mathfrak{o}_v$ intersect in the special fibre.

Let $\beta : U \to \mathbb{P}^1 - \{0, 1, \infty\}$ be a finite map defined on an open $U \subseteq X$. We lift s to a section of $\pi_1(U/k)$ and apply Theorem 18 to this lift and $\beta^* E_\lambda/U$. Since $\underline{a}(s)$ does not change with the lift, we are still in the second case of the conclusion of Theorem 18. The 2-adic representation induced by the section factors through

$$\begin{pmatrix} \delta\varepsilon & * \\ & \delta \end{pmatrix} \subseteq \mathrm{GL}_2(\mathbb{Z}_2),$$

which is a pro-2 group. Now by passing to a neighbourhood $h : U' \to U$ of s we may assume that

$$\rho_{h^*\beta^*E_\lambda/U',2} : \pi_1(U') \to \mathrm{GL}_2(\mathbb{Z}_2)$$

factors through a pro-2 group. Let S be a finite set of places containing all the bad places for $h^*\beta^*E_\lambda/U'$ and the places dividing 2. Let $\mathscr{U}'/\operatorname{Spec}(\mathfrak{o}_{k,S})$ be a hyperbolic curve model of U'/k. Denote by $\pi_1^{(2)}(-)$ the fibrewise pro-2 fundamental group. Then the representations factor as

$$\begin{tikzcd}
\pi_1^{(2)}(U') \arrow[r, two heads, "\mathrm{sp}"] \arrow[d, shift right, "\mathrm{pr}_*"'] & \pi_1^{(2)}(\mathscr{U}') \arrow[r, "\rho_{h^*\beta^*E_\lambda/U',2}"] \arrow[d, "\mathrm{pr}_*"'] & \mathrm{GL}_2(\mathbb{Z}_2) \\
\mathrm{Gal}_k \arrow[u, shift right, "s"'] \arrow[ur, "\mathrm{sp}\circ s"] \arrow[r, two heads] & \pi_1(\operatorname{Spec}(\mathfrak{o}_{k,S})) &
\end{tikzcd}$$

The specialisation map $\mathrm{sp} : \pi_1^{(2)}(U') \to \pi_1^{(2)}(\mathscr{U}')$ is an isomorphism on the kernels of the respective projections pr_*. Let $v \nmid 2$ be a cuspidal or degenerate place, and let $D_v \subset \mathrm{Gal}_k$ be a choice of a decomposition subgroup at v. Then, as a_v degenerates into y_v we find that

$$\mathrm{sp} \circ s(D_v) \text{ cyclotomically normalizes } I_{y'_v} \tag{6.1}$$

where $I_{y'}$ denotes the inertia group of $y' \in Y' = \overline{X'} \setminus X'$ in $\pi_1^{(2)}(U'_{\bar{k}}) \subseteq \pi_1^{(2)}(\mathscr{U}')$. Here **cyclotomically normalizing** means that the induced action by conjugation is via the cyclotomic character. Due to neglecting base points and choice of prolongations of places, these inertia and decomposition groups are only well defined up to conjugation and (6.1) has to be considered as holding for suitable choices within the conjugacy classes of these groups. Because of

$$\rho_{s,h^*\beta^*E_\lambda/U',2}(D_v) \subseteq \begin{pmatrix} \delta\varepsilon & * \\ & \delta \end{pmatrix},$$

depending on whetheror not $* = 0$ for the restriciton of the representation to D_v, there are two, namely

$$\begin{pmatrix} 1 & * \\ & 1 \end{pmatrix} \text{ and } \begin{pmatrix} 1 & \\ * & 1 \end{pmatrix},$$

or a unique maximal unipotent subgroup normalized by D_v. But in any case only

$$\begin{pmatrix} 1 & * \\ & 1 \end{pmatrix}$$

is cyclotomically normalized. This maximal unipotent subgroup is independent of v and thus must by (6.1) be conjugate to the image of $I_{y'_v}$. We conclude with Lemma 27 that

$$\beta(y_v) = \beta(h(y'_v)) \in \{0, 1, \infty\}$$

is also independent of v.

By Riemann-Roch, it is easy to find for any partition $Y = Y_0 \amalg Y_\infty$ a suitable $U \subseteq X$ and a finite $\beta : U \to \mathbb{P}^1 - \{0,1,\infty\}$ with $\beta(Y_0) = \{0\}$ and $\beta(Y_\infty) = \infty$. We conclude that $v \mapsto y_v$ must be a constant function, i.e., all local points a_v degenerate or are cuspidal with the very same point $y \in Y$. This means in particular, that the residue field extension $\kappa(y)/k$ has a split place above almost all places v of k, and this is only possible if $\kappa(y) = k$ by Lemma 30 below. Thus $y \in Y(k)$ and this finishes the proof. $\square$

Remark 29. Note that the use of the limit argument in Theorem 28 does again work without the use of Faltings' Theorem on the finiteness of the set of rational points for curves of genus at least 2 over number fields.

Lemma 30. *Let F/K be a finite extension of number fields such that for all but finitely many places of v there is a place w of E with the same residue field. Then we have necessarily $E = F$.*

Proof: Let E/K be a Galois hull of F/K and let $G = \mathrm{Gal}(E/K) \supseteq H = \mathrm{Gal}(F/K)$ be the respective Galois groups. The assumption says that for all but finitely many v the conjugacy class of Frobenius elements at places $w \mid v$ meets H nontrivially. But since every element of G is a Frobenius element infinitely often, this implies that G is the union of the conjugates of H. This is only possible if $G = H$ and thus $F = K$ as claimed. $\square$

7. On the birational section conjecture over $\mathbb{Q}$ with conditions

In the case of hyperbolic curves over $\mathbb{Q}$, we offer the following version of a birational section conjecture with conditions.

Theorem 31. *Let $X/\mathbb{Q}$ be a hyperbolic curve. A birationally liftable section $s : \mathrm{Gal}_\mathbb{Q} \to \pi_1(X)$*

(a) *comes from a rational point $a \in X(k)$,*

(b) *or is cuspidal,*

if and only if for every open $U \subseteq X$ with geometric point $\bar{u} \in U$, a lift $\tilde{s} : \mathrm{Gal}_\mathbb{Q} \to \pi_1(U, \bar{u})$, and every family E/U of elliptic curves with geometric fibre $E_{\bar{u}}$ over $\bar{u}$ the associated family of 2-dimensional representations

$$\big(\rho_{\tilde{s},E/U,\ell} \ : \ \mathrm{Gal}_\mathbb{Q} \to \mathrm{GL}(\mathrm{T}_\ell\, E_{\bar{u}})\big)_\ell$$

has the following properties

(c) *either there exists a finite set of places S independent of ℓ such that $\rho_{\tilde{s},E/U,\ell}$ is unramified outside ℓ and the places in S,*

(d) *or there is a quadratic character $\delta : \mathrm{Gal}_\mathbb{Q} \to \{\pm 1\}$ such that for all ℓ we have an exact sequence*

$$0 \to \delta\varepsilon \to \rho_{\tilde{s},E/U,\ell} \to \delta \to 0$$

where ε is the ℓ-adic cyclotomic character.

Remark 32. Assumption (c) rules out that infinitely many places of k are degenerate with respect to the adele associated to the birationally liftable $s : \mathrm{Gal}_\mathbb{Q} \to \pi_1(X)$ and $X \subseteq \overline{X}$. But it does not exclude that maybe infinitely many places are cuspidal.

Proof: Step 1: Let $\overline{X}$ be the smooth completion of X. We show first that (a) or (b) implies (c) or (d). Let $s : \mathrm{Gal}_\mathbb{Q} \to \pi_1(X)$ be a birationally liftable section and let E/U be a family of elliptic curves and $\tilde{s} : \mathrm{Gal}_\mathbb{Q} \to \pi_1(U)$ as in the theorem. Upon replacing U by neighbourhood of $\tilde{s}$ we may assume that E/U has either good or bad semistable reduction along points in $\overline{X} \setminus U$. If the section s originates from a rational point or a cusp located in the point $a \in \overline{X}(k)$, then the representation $\rho_{s,E/U,\ell}$ has property (c) if the family has good reduction in a, because by functoriality then $\rho_{\tilde{s},E/U,\ell} = \rho_{E_a,\ell}$ is the representation of the corresponding fibre of (the extended) family in a. If the reduction is bad semistable, then $\rho_{\tilde{s},E/U,\ell}$ has property (d).

Step 2: Now we assume (c) or (d) holds for any family E/U as in the theorem. By Theorem 28 and Theorem 18 we are done if (d) applies. We therefore assume that (c) holds for a given family E/U with bad reduction along $\overline{X} \setminus U$, and that S is a finite set of places as in (c).

Step 3: Let $\bar{\rho}_{E/U,\ell}$ be the mod ℓ reduction of $\rho_{E/U,\ell} : \pi_1(U) \to \mathrm{GL}(\mathrm{T}_\ell E_{\bar{u}})$. If the mod ℓ represenation $\bar{\rho}_\ell$ of $\rho_{\tilde{s},E/U,\ell}$ is reducible, then we can pass to the neighbourhood $h : U' \to U$ defined by

$$(\bar{\rho}_{E/U,\ell})^{-1}(\operatorname{im} \bar{\rho}_\ell) \subseteq \pi_1(U).$$

The pull back family h^*E/U' carries by construction a family of cyclic subgroups of degree ℓ, hence provides a non-constant map

$$U' \to \mathscr{E}\ell\ell_0(\ell)$$

where $\mathscr{E}\ell\ell_0(\ell)$ is the corresponding modular curve over $\mathbb{Q}$. If $\ell = 11$ or $\ell \geq 17$ then the Jacobian of $\mathscr{E}\ell\ell_0(\ell)$ posseses quotient abelian varieties A with $A(\mathbb{Q})$ and $\text{Ш}(A/\mathbb{Q})$ finite, see Example 7. We conclude by Stoll's theorem as recalled in Corollary 15 that the image of $\underline{a}(s)$ in $\mathscr{E}\ell\ell_0(\ell)$ has finite support, whence by Corollary 9 we may deduce that s is either cuspidal or s_a for some $a \in X(k)$.

Step 4: We may therefore assume that $\bar{\rho}_\ell$ is irreducible for $\ell = 11$ or $\ell \geq 17$. Since $\det(\rho_{s,E/U,\ell}) = \varepsilon$ is cyclotomic, the representation $\bar{\rho}_\ell$ is odd. Then complex conjugation $c \in \mathrm{Gal}_{\mathbb{Q}}$ acts like

$$\bar{\rho}_\ell(c) \sim \begin{pmatrix} 1 & \\ & -1 \end{pmatrix}$$

so that any invariant subspace of dimension 1 after extension of scalars must be an eigenspace of $\bar{\rho}_\ell(c)$ and thus already defined over $\mathbb{F}_\ell$. Hence $\bar{\rho}_\ell$ is absolutely irreducible for $\ell = 11$ or $\ell \geq 17$. By Serre's modularity conjecture proven by Khare and Wintenberger [KW09] Theorem 1.2, the representation $\bar{\rho}_\ell$ is modular, and even of the predicted level and weight.

Since locally s comes from a point we see that $\bar{\rho}_\ell$ is finite locally at ℓ. We deduce from [Se87] Proposition 4, that the predicted weight is $k = 2$. Concerning the level we restrict[1] to primes $\ell = 11$ or $\ell \geq 17$ subject to the following congruences

(i) $\ell \equiv 3 \mod 8$,
(ii) $\ell \equiv 2, 5 \mod 9$,
(iii) $\ell \not\equiv \pm 1 \mod p$ for all $p \in S \setminus \{2, 3\}$.

This bounds the order p^c of the p-Sylow subgroups for $p \neq \ell$ and $p \in S \cup \{2, 3\}$ in $\mathrm{GL}_2(\mathbb{F}_\ell)$, which is of order $(\ell - 1)^2 \ell (\ell + 1)$ to

(i) $p^c \leq 2^4$ for $p = 2$,
(ii) $p^c \leq 3$ for $p = 3$,
(iii) and $p^c = 1$ for $p \in S \setminus \{2, 3\}$.

By [Se87] Proposition 9, such bounds on the wild inertia yield the following bound for the predicted conductor: it must be a divisor of

$$N = 2^{12} \cdot 3^5 \cdot \prod_{p \in S \setminus \{2,3\}} p^2.$$

It follows that for every such ℓ there is an eigenform of weight $k = 2$ and level a divisor of N such that the corresponding Galois representation modulo ℓ agrees with $\bar{\rho}_\ell$. Since there are only finitely many such eigenforms, we can pick one f with ℓ-adic representation $\rho_{f,\ell}$ that occurs for infinitely many ℓ. Then we obtain congruence for traces of Frobenius for all $p \notin S$ (unramified in the respective mod ℓ representation)

$$a_p(f) \equiv \mathrm{tr}(\mathrm{Frob}_p | \rho_{f,\ell}) \equiv \mathrm{tr}(\mathrm{Frob}_p | \bar{\rho}_\ell) \equiv a_p(\rho) \mod \ell.$$

---

[1] It is time to acknowledge the influence of the paper [HV11] by Helm and Voloch. The argument to follow goes back essentially to Serre [Se87].

Since both sides are integral over $\mathbb{Z}$ and independent of ℓ we conclude that

$$a_p(f) = a_p(\rho) \in \mathbb{Z}$$

as integers. It follows that the abelian variety quotient $J_0(N) \to E_f$ associated to f is of dimension 1, an elliptic curve, and that

$$\rho_{E_f,\ell} \otimes \mathbb{Q}_\ell = \rho_{f,\ell} \otimes \mathbb{Q}_\ell \cong \rho_{s,E/U,\ell} \otimes \mathbb{Q}_\ell. \tag{7.1}$$

This allows us to read off weights of Frobenius: an elliptic curve over $\mathbb{Q}$ has good reduction outside a finite set of places, the corresponding local representations are unramified and of weight -1, and thus all but finitely many places are integral with respect to $\underline{a}(s)$ and $X \subseteq \overline{X}$. This almost shows that $\underline{a}(s) \in U(\mathbb{A}_\mathbb{Q})_\bullet$ with the exception of finitely many places p where

$$a_p \in Y(\mathbb{Q}_p) = (\overline{X} \setminus U)(\mathbb{Q}_p).$$

(We sincerely hope that the reader is not confused by the two meanings of a_p: as a local point associated to the section, and as a trace of Frobenius.)

We may assume that the isomorphism in (7.1) respects the two integral structures to the extend that $\rho_{s,E/U,\ell}$ acts on a lattice in $\mathrm{T}_\ell\, E_f$ with cyclic quotient. As $\bar{\rho}_\ell$ is irreducible for $\ell \gg 0$, this cyclic quotient can be nontrivial only for finitely many ℓ, whence we can replace E_f by an isogenous elliptic curve E'_f and obtain even isomorphisms on the level of $\mathbb{Z}_\ell$-coefficients

$$\rho_{E'_f,\ell} \cong \rho_{s,E/U,\ell}. \tag{7.2}$$

Step 5: Let p be a place such that $a_p = y \in Y(\mathbb{Q}_p)$. Then the local section $s_p = s \otimes \mathbb{Q}_p$ is cuspidal and $\rho_{s,E/U,\ell}|_{\mathrm{Gal}_{\mathbb{Q}_p}}$ factors through a decomposition group at y, hence through a group

$$\begin{pmatrix} \delta\varepsilon & * \\ & \delta \end{pmatrix}$$

with δ a quadratic character. Thus the semisimplification of the local representation has Frobenius weights $-2, 0$. It follows that E'_f cannot have potentially good reduction at p since otherwise the local Frobenius weights at p were -1. By passing to a finite extension $K/\mathbb{Q}_p$ we may assume that E'_f has split semistable reduction. Let $q \in K^*$ be the local parameter in the description as a Tate elliptic curve. Then $\rho_{E'_f,\ell}|_{\mathrm{Gal}_K}$ as an extension is described by

$$q \in K^* \subset \widehat{K^*} = \mathrm{Ext}^1_{\mathrm{Gal}_K}(\mathbf{1}, \varepsilon),$$

and this agrees with the extension class

$$[\rho_{s,E/U,\ell}|_{\mathrm{Gal}_K}] = q \in \mathrm{Ext}^1_{\mathrm{Gal}_K}(\mathbf{1}, \varepsilon),$$

described by $\rho_{s,E/U,\ell}|_{\mathrm{Gal}_K}$.

Let D_y be the decomposition subgroup at the cusp y in $\pi_1(U \otimes K)$ and let $\hat{\mathbb{Z}}(1) = I_y \subset D_y$ be the inertia subgroup. The section s being cuspidal at p, we find the following diagram.

$$\begin{tikzcd}
1 \arrow[r] & \hat{\mathbb{Z}}(1) \arrow[r] \arrow[d, "m\cdot"] & D_y \arrow[r] \arrow[d, "\rho_{E/U,\ell}|_{D_y}"] & \mathrm{Gal}_K \arrow[r] \arrow[d, "\varepsilon"] \arrow[l, bend right, "s_p"'] & 1 \\
1 \arrow[r] & \begin{pmatrix} 1 & * \\ & 1 \end{pmatrix} \arrow[r] & \begin{pmatrix} \varepsilon & * \\ & 1 \end{pmatrix} \arrow[r] & \mathbb{Z}_\ell^* \arrow[r] & 1
\end{tikzcd}$$

with $m \in \mathbb{Z}$ related to the valuation of the Tate parameter governing the semistable reduction of E/U in y. Choosing a parameter t at y we can split $1 \to \hat{\mathbb{Z}}(1) \to D_y \to \mathrm{Gal}_K \to 1$ with a section s_t using the corresponding tangential base point. The difference of s_p and s_t yields a class

$$\delta(s_p,s_t) \in \mathrm{H}^1(K,\hat{\mathbb{Z}}(1)) = \mathrm{Ext}^1_{\mathrm{Gal}_K}(\mathbf{1},\varepsilon),$$

which is the class of the cocycle $\sigma \mapsto s_p(\sigma)/s_t(\sigma) \in \hat{\mathbb{Z}}(1)$. We conclude that

$$q = [\rho_{s,E/U,\ell}|_{\mathrm{Gal}_K}] = [\rho_{E/U,\ell}|_{D_y} \circ s_t] + m\delta(s_p,s_t) \in \mathrm{Ext}^1_{\mathrm{Gal}_K}(\mathbf{1},\varepsilon).$$

If $Q(t) \in K((t))$ is the local Tate parameter governing the semistable reduction of E/U in y, then

$$[\rho_{E/U,\ell}|_{D_y} \circ s_t] = s_t^*(Q(t)) \in K^* \subset \widehat{K^*} = \mathrm{H}^1(K,\hat{\mathbb{Z}}(1))$$

because the pullback with $s_t : \mathrm{Gal}_K \to \mathrm{Gal}_{K((t))} = D_y$

$$s_t^* : \mathrm{H}^1(K((t)),\hat{\mathbb{Z}}(1)) \to \mathrm{H}^1(K,\hat{\mathbb{Z}}(1))$$

maps $K \cdot t^{\mathbb{Z}} = \mathrm{im}\big(K((t))^* \to \widehat{K((t))^*}\big)$ to $K^* \subset \widehat{K^*}$ by evaluating at $t=1$. Consequently,

$$m\delta(s_p,s_t) \in K^* \subset \mathrm{H}^1(K,\hat{\mathbb{Z}}(1))$$

and since $\widehat{K^*}/K^* = \hat{\mathbb{Z}}/\mathbb{Z}$ is uniquely divisible, also

$$\delta(s_p,s_t) \in K^* \subset \mathrm{H}^1(K,\hat{\mathbb{Z}}(1)).$$

We deduce that s_p is also tangential, i.e., can be constructed from a tangent vector at y.

Step 6: We may assume that U carries a non-constant unit $u : U \to \mathbb{G}_m$. Because either $a_p \in U(\mathbb{Q}_p)$ with even integral coefficients for all but finitely many p, or s_p is tangential, we find that $u(s) = u_*(s)$ yields an adelic value in $\mathbb{G}_m(\mathbb{A}_{\mathbb{Q}})$ that survives all finite descent obstructions. By Proposition 4, we deduce that $u(s) \in \mathbb{Q}^*$. It follows that s has finite support in

$$u^{-1}(u(s)) \amalg Y \subseteq \overline{X}.$$

By Corollary 9 the section s belongs to a rational point of U. This completes the proof. ☐

References

[EsHa08] Esnault, H., Hai, Ph. H., Packets in Grothendieck's Section Conjecture, *Advances in Mathematics* **218** (2008), no. 2, 395–416.

[Ha08] Harari, D., Le défaut d'approximation forte pour les groupes algébriques commutatifs, *Algebra Number Theory* **2** (2008), no. 5, 595–611.

[HSx10] Harari, D., Stix, J., Descent obstruction and fundamental exact sequence, in: *The arithmetic of fundamental groups — PIA 2010*, editor Jakob Stix, Contributions in Mathematical and Computational Science **2**, Springer, 2012, chapter 7, 147–166.

[Gr83] Grothendieck, A., Brief an Faltings (27/06/1983), in: *Geometric Galois Action 1* (ed. L. Schneps, P. Lochak), LMS Lecture Notes **242**, Cambridge, 1997, 49–58.

[HV11] Helm, D., Voloch, J. F., Finite descent obstruction on curves and modularity, *Bull. Lond. Math. Soc.* **43** (2011), no. 4, 805–810.

[KW09] Khare, C., Wintenberger, J.-P., Serre's modularity conjecture I, *Invent. Math.* **178** (2009), no. 3, 485–504.

[KL89] Kolyvagin, V. A., Logachëv, D. Yu., Finiteness of the Shafarevich-Tate group and the group of rational points for some modular abelian varieties. (Russian) *Algebra i Analiz* **1** (1989), no. 5, 171–196; translation in *Leningrad Math. J.* **1** (1990), no. 5, 1229–1253.

[KL92] Kolyvagin, V. A., Logachëv, D. Yu., Finiteness of over totally real fields, *Math. USSR Izvestiya* **39** (1992), no. 1, 829–853.

[Ko05] Koenigsmann, J., On the 'section conjecture' in anabelian geometry, *J. Reine Angew. Math.* **588** (2005), 221–235.

[La65] Lazard, M., Groupes analytiques p-adiques, *Publications mathématiques de l'IHES* **26** (1965), 5–219.

[Mz04] Mochizuki, S., Noncritical Belyi maps, *Math. J. Okayama Univ.* **46** (2004), 105–113.

[Na90] Nakamura, H., Galois rigidity of the étale fundamental groups of punctured projective lines, *J. Reine Angew. Math.* **411** (1990), 205–216.

[NSW08] Neukirch, J., Schmidt, A., Wingberg, K., *Cohomology of number fields*, second edition, Grundlehren der Mathematischen Wissenschaften **323**, Springer, 2008, xvi+825 pp.

[Se87] Serre, J.-P., Sur les représentations modulaires de degré 2 de Gal($\overline{\mathbb{Q}}/\mathbb{Q}$), *Duke Math. Journal* **54** (1987), no. 1, 179–230.

[St07] Stoll, M., Finite descent obstructions and rational points on curves, *Algebra & Number Theory* **1** (2007), 349–391.

[Sx11] Stix, J., *Evidence for the section conjecture in the theory of arithmetic fundamental groups*, Habilitationsschrift, Heidelberg, 2011, x+190pp, to appear as Springer Lecture Notes in Mathematics.

[Sx12] Stix, J., Birational p-adic Galois sections in higher dimension, arXiv:1202.2781v1[math.AG].

Jakob Stix, MATCH - Mathematisches Institut, Universität Heidelberg, Im Neuenheimer Feld 288, 69120 Heidelberg, Germany
E-mail address: `stix@mathi.uni-heidelberg.de`
URL: `http://www.mathi.uni-heidelberg.de/~stix/`